

California Coastal Salmon and Steelhead Current Stream Habitat Distribution Table

Table to be used with DeLorme Topo Quads, Bibliography, and Contacts & Expertise list

(Streams listed from north to south)

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
<i>Chinook Salmon - California Coastal ESU</i>					
<i>Coho Salmon - Central California Coast ESU</i>					
<i>Steelhead - Northern California Coast Steelhead ESU</i>					
Mendocino County streams					
Whale Gulch 24N 19W, Sec. 4	SH	RM-0.3	Jones	Personal observations: the barrier is a falls some 25 feet high.	1968
	COH	RM-?	Jones	Electrofishing: 2 stations, one below the barrier and one above, found juvenile SH below and RT above in 1984 .	1984
	RT	RM-?	Brown and Moyle	Report: COH reported in this stream; "Status of Coho Salmon in California", 1991	1991
Low Gap Creek 24N, 19W, Sec. 9	None	Barrier @ RM-0	Jones	Personal observations: barrier near confluence with the ocean.	1984
Flat Rock Creek 24Nm 19W, Sec. 9	None	Barrier @ RM-0	Jones	Personal observations: there is a barrier near the confluence with the ocean.	1984
Bear Harbor Creek 24N, 19W, Sec. 26	SH	Mainstem-RM-0.4	Ginger & Jones	Personal observations: there is a barrier in the upper Middle Fork 8 feet High. Electrofished: survey found juvenile SH present.	1979
		NF- RM-0.1 MF- RM-0.7 SF- 0.06	Jones		1984

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Jackass Creek T23N, R19W, Sec. 1	COH	RM-?	Murphy	COH and SH were present.	1948
	SH	RM-1.5	Grass	Personal observation: juvenile SH were present.	1963-1970
			Jones	Personal observations: good habitat for SH was found. Electrofishing in 1983 & 1987 found juvenile SH.	1983 1987
			Griffin, Salomone & Long	Stream Survey: juvenile SH were present.	1988
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
North Fork Jackass Creek T23N, R18W, Sec.6	SH	RM-2	Jones	Survey: Good habitat was found in the North Fork for salmonids. Appears ideal habitat for COH but found only juvenile SH.	1983
			Harris	Survey: faceplate snorkel survey found only juvenile SH. They were present in multiple year classes.	1998
East Branch North Fork Jackass Creek	SH	RM-?	Jones	Personal observations: juvenile SH were present in the mouth of the stream.	1983
Little Jackass Cr. T23N, R18W, Sec. 7	None	RM-0	Jones	Electrofished: but found no fish. This stream appears to have good fish habitat.	1985

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Usal Creek 23N, 18W, Sec. 22	SH	RM-6.8	Shapovalov	Fish rescue report: reported 3,963 COH juveniles were collected in 1940.	1940
	COH	RM-?	Shapovalov	Fish rescue report: reported collecting 61,133 in 1945.	1945
			Kimsey	Fish rescue report: reported collecting 13,864 in 1952.	1953
			Cherr & Griffin	Stream inventory: Cherr & Griffin reported SH present in the drainage.	1979
			Moore	Memo: Moore reported to CDF, a barrier some 50 feet high, to anadromous fish life.	1981
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Coyle, Mesman, Hickethier & Young	Electrofished: 9 sites, found juvenile SH 36,011 feet above the mouth.	1995
South Fork Usal Creek 23N, 18W, Sec.22	COH	up to Julias Creek	Robert	Electrofished: juvenile COH were present, (only one year class of the three was found).	1987, 1993
	SH	RM-1.5 RM-3.0	Cherr & Griffin	SH were present.	1979
			Jones	Electrofished: found juvenile SH.	1987
			Mesman & Hickethier	Electrofished: captured only juvenile SH, young of year age size, yearling age size, and yearling + age size fish. Electrofished: 2 sites, found juvenile SH 9,194 feet upstream.	1993 1995?
Julias Creek T23N, 18W, Sec.24	SH	RM-1.0	Jones	Electrofished: caught juvenile SH; young of the year and yearling + age sized fish.	1987
			Mesman, & Young	Electrofished: found juvenile SH 4, 237 feet upstream.	1995
Soldier Creek T23N, R18W, Sec. 14	SH	RM-1.7	Mesman & Young	Electrofished: 2 sites, found juvenile SH up to 8,945 feet upstream.	1995

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Little Bear Creek T23N, R18W, Sec. 11	SH	RM- 243 feet	Mesman & Young	Electrofished: 1 site, found juvenile SH 243 feet upstream.	1995
Bear Creek T23N, R18W, Sec. 11	SH	RM-559 feet	Mesman & Young	Electrofished: 1 site, found juvenile SH 559 feet upstream.	1995
Chimney Rock Creek T23N, R18W, Sec. 4	SH	RM-1.0	Coyle & Young	Electrofished: 1 site, found juvenile SH present.	1995
Cottaneva Creek 22N, 18W, Sec, 23	COH	RM-6.0	Jone	Electrofished: found juvenile SH below Middle Fork, but no juvenile COH caught in 1983.	1983
	SH		Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Jones	Electrofished: a ratio of 240 juvenile SH found for every COH.	1997
			Harris	Carcass survey: Dec 1998, found no adult fish in a one mile section of stream near the LP Demonstration Forest area.	1998
			Harris	Carcass survey: Between the Middle Fork and Rockport. No fish were seen, the area was heavily impacted by cattle in Dec.	1999
North Fork Cottaneva Creek 22N, 18W, Sec. 2	COH	RM-1.3	Edie	Survey: juvenile COH and SH were observed. Good potential for spawning, little value as a nursery area.	1966
	SH		Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Jones	Good nursery habitat below Dunn Creek.	1997
			Coyle & Young	Electrofished: One site, found juvenile SH within 459 feet of Dunn Creek. No COH were present.	1995
			Daugherty	Electrofished: caught juvenile COH above Dunn Creek, none in the station shocked in mid section 36.	1996 & 1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Dunn Creek 22N, 18W, Sec 1	COH SH	RM-0.6 RM-1.5	Mesman & Young Grass Harris	Electrofished: 3 sites, juvenile SH up to 5,682 feet upstream. Personal observation: juvenile COH were present below the Hwy 1 culvert, which was currently a barrier to them. CDFG has requested Cal trans to modify this hwy culvert. SH adults have been seen upstream to about 1/4 mi. above the two forks and into the unnamed fork to the east. Personal observation: still having problems getting Cal Trans to fix the culvert for fish passage.	1995 1999 2000
Middle Fork Cottaneva Creek 22N, 18W, Sec. 2	COH SH	RM-? RM-1.0	Edie Coyle & Young Daugherty	Survey: "Good spawning gravels near the upper end of the fisheries value, many 2 year old steelhead. The lower portions have almost no 2 year old fish. Juvenile COH were observed in the lower portions." (Edie did not note the upper limit of COH). Electrofished: One site, juvenile SH were present 1,500 feet upstream. Personal observation: L.P. found COH in the mouth, but not in Sec. 12. SH were present in both areas. "Good spawning gravels near the upper end of fisheries value, containing many 2 year old juvenile steelhead. The lower portions have almost no 2 year old fish."	1966 1995 1966
Powder House Gulch T22N, R18W, Sec. 13	SH	RM-0.7	Daugherty	Electrofished: Found SH present.	1994 & 1995
South Fork Cottaneva Creek 22N, 18W, Sec. 24	COH SH	RM-2.5 RM-2.5	Santos & Crowdus Edie Brown and Moyle Daugherty Mesman & Young	Survey: Santos and Crowdus saw no juvenile COH and SH during their survey. Survey: Edie saw juvenile COH and SH during his survey. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: Daugherty found juvenile COH, all three years. Electrofished: 3 sites, juvenile COH were found up to 7,745 feet and juvenile SH were present up to 11,480 feet.	1961 1966 1991 1994, 1995, 1996 1995

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Slaughterhouse Gulch T22N, R17W, Sec. 19	COH	RM-0.1	Daugherty	Electrofished: juvenile COH were present but only near the mouth.	1994, 95, and 96
	SH	RM-1.0	Mesman & Young Jones & Grass	Electrofished: 2 stations, Juvenile SH were found 4,422 feet upstream. Electrofished: juvenile SH were present.	1995 1999
Kimball Gulch T22N, R17W, Sec. 19	COH	RM-?	Jones & Grass	Electrofished: Juvenile COH were present but the extent of occupancy was unknown.	1995
	SH	RM-?	Daugherty	Electrofished: Juvenile SH were present near the mouth, but the extent of occupancy was unknown.	1999
Rockport Creek 22N, 18W, Sec.25	COH	RM-0.35	Edie	Survey: juvenile COH are present but the extent of occupancy was unknown.	1966,
	SH	RM-0.5	Mesman & Young	Electrofished: one site, juvenile SH were found 1,843 feet upstream.	1995
			Daugherty Grass	Electrofished: juvenile SH were present but the extent of occupancy was unknown. Personal observations: juvenile SH were present.	1966, 1999
Hardy Creek 21N, 18W, Sec. 1	COH	RM- 1.7	Dieden & Jones	Juvenile COH and SH were present up to the barrier at 1.7 miles.	1961
	SH	RM-1.7	Edie	Survey: a logging road barrier 10 feet high was found above Bear Creek.	1966
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Daugherty	Electrofished: Daugherty found juvenile SH above the barrier.	1994, 1995, 1996
			Cooper & Mesman	Electrofished: 2 sites, juvenile juvenile SH were present up to the confluence of the Middle and North Forks.	1997
Jones & Grass	Personal observation: no COH were present in this drainage.	1999			

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
North Fork Hardy Creek 22N, 17W, Sec. 32	SH	Above Middle Fork (RM-0.04)	Santos & Crowdus	Survey: juvenile SH present, the drainage becomes steep 75 yards above the Middle Fork.	1961
			Flosi & Long	Survey: a barrier of 9 feet in height was present 195 feet upstream.	1986
South Fork Hardy Creek or Bear Creek T22N, R17W, Sec. 22	SH	RM-0.2	Cooper & Mesman	Electrofished: found juvenile steelhead 896 feet upstream.	1997
Middle Fork Hardy Creek 22N, 17W, Sec.29	SH	RM-0.1	Nye & Dieden	Survey: the juvenile SH were considered scarce and were present only for the first 175-200 yards. They numbered one to 10 in the larger pools.	1961
			Retherford & Lester	Electrofished: found juvenile SH 275 feet upstream.	1998
Juan Creek 21N, 18W, Sec 1.	SH	RM-3.3	Jones	Survey: juvenile SH were present in 1961. No juvenile COH were seen during the survey.	1961
	COH	RM-4.2	Primbs	Survey: 45 juvenile COH and 187 juvenile SH were netted and examined in 1966.	1966
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Daugherty	Electrofished: juvenile SH were present.	1996
			Grass and Jones	Personal observation: no juvenile COH currently present in Juan Creek.	1996- 1999

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Little Juan Creek 21N, 17W, Sec 6	SH COH	RM-2.0 RM-?	Crowdus & Santos Primbs Brown and Moyle Humphrey Daugherty Jones	Survey: juvenile SH were found in 1961. Survey: found 57 juvenile SH. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Personal information: 5 redds seen (species unknown). Electrofished: found juvenile SH and COH. Personal observation: COH have not been seen in this drainage recently (W. Jones).	1961 1966 1991 1993 1996 1997
Little Howard Cr. 21N, 17W, Sec. 7	None	RM-0	Jones	Survey: A barrier was present at the mouth, (Cliff) and the Hwy 1 culvert maintains it. However, the stream habitat looks suitable for salmonids.	1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Howard Creek 21N, 17W, Sec. 18	SH	RM-2.6	Crowdus & Dieden	Survey: juvenile SH were present, but no COH.	1966
	COH	RM-2.0 +?	Thomas & Primbs	Survey: caught 155 juvenile SH, no COH.	1966
			Grass	Electrofished: juvenile SH were present but no COH.	1983
			Jones	Electrofished: caught juvenile SH, no COH.	1986
			Jones	Electrofished: COH caught in 1988 were juveniles from hatchery plants in early 1988. No COH in 1989-91.	1988-91
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Harris	1995 survey: juvenile SH were present but no COH were seen.	1995,
			Poe	Planting receipt: --yearling COH were planted to re-establish the full three year cycle.	1996-96-97
		Harris	Electrofishing: found juvenile COH and juvenile SH. The COH were from hatchery plants.	1998	
		Harris	Electrofishing: caught juvenile COH and SH.	1999	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
De Haven Creek 21N, 17W, Sec. 18	SH	RM-2.9	Nye & Santos	Surveys: found juvenile COH & SH present; 1-15 COH/pool and 20 juvenile SH/ pool. Also found juvenile SH above Martin Creek tributary (North Fork).	1961
	COH	RM-1.0	Jones & Dieden	Survey: found SH young of the year age and yearling age sized fish.	1961
			Primbs	Survey: juvenile SH were seen to 1,900 feet upstream.	1966
			Brown & Brown	Electrofished: caught juvenile SH.	1988-91
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Jones	Electrofished: in the lower portion of GP land, caught juvenile SH.	1995
			GP	Electrofished: sampled two locations in 1998, juvenile SH were found in both the lower and upper area.	1988
			Amado & Emig	Personal information: CDFG planted yearling COH in the spring of 1988.	1988
			Jones	Electrofished and found juvenile COH in November.	1988
			CDFG	CDFG planted yearling COH along the access road in the lower area in 1995, 1996, and 1997.	1995-97
			Ludtke	Electrofished: no COH present, juvenile SH were present.	1996
		Harris	Electrofished: no COH present, juvenile SH were found. Electrofished: only found juvenile SH.	1998 1999	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Wages Creek 21N, 17W, Sec. 30	COH	RM- 3.6+	Edie	Survey: no juvenile COH were observed in the main stem of Wages Creek in 1966. Juvenile SH were very abundant in most of the stream, possibly due to the amount of riffle.	1966
	SH	RM-7.2	Amado	Personal observation: yearling COH were planted in the stream.	1988
			Jones	Electrofished: caught juvenile COH and SH in the spring and fall of 1988. Found juvenile SH present 1989, 1990, 1991.	1988-89 & 91
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			GP	Electrofished: 1993-95, and 1998 (including lower GP area in 1995 and '98), found juvenile SH each year. Caught juvenile COH, in the lower GP area in the fall of 1995 and in upper stream in 1998. No COH were caught in 1993 and 1994.	1993-95 1998
			Jones	Personal observation: CDFG planted yearling COH in 1995-96-97.	1997
			Harris	Carcass survey: found two COH grilse.	1999
			Harris	Downstream migrant trap operated in the spring, found juvenile COH and SH.	1999-00
North Fork Wages Creek 21N, 17W, Sec. 26	SH	RM- 0.5	Edie	Survey: North Fork, juvenile SH were present, good spawning potential, gravel present in fair quantities. Little nursery value due to small flows.	1966
South Fork Wages Creek 21N, 16W, Sec. 31	SH	RM- 0.3	Edie	Survey: South Fork, juvenile SH were present, good spawning potential, gravel present in fair quantities. Little nursery value due to small flows.	1966
Rider Gulch 21N, 17W, Sec. 29	SH	RM- 1.0	Edie	Survey: juvenile SH and COH were observed in Rider Creek. Fair spawning area, some gravel, almost no nursery value, very low flow in July.	1966
	COH	RM- 1.0			
Chadbourne Gulch 20N, 17W, Sec. 8	SH	RM- 1.0	Grass & Aasen	Electrofished: collected young-of-the-year age and yearling age sized SH.	1983

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³	
Abalobadiah Creek 20N, 17W, Sec. 28	SH	RM- 2.6 Plus "middle" and "south" forks	Santos	Survey : in 1961 included unnamed tributaries, (middle and south forks). The juvenile SH density averaged one to 8 fish per pool in the Middle Fork and one to 6 fish per pool in the South Fork. No COH were observed.	1961	
Seaside Creek 20N, 17W, Sec. 33	SH	RM- 1.0	Cherr & Griffin	Survey: juvenile SH and COH were present in 1979.	1979	
	COH	RM- 1.0	Jones	Personal observation: Lower area, dry to intermittent in the summer. No COH were present.	1997	
Ten Mile River 20N, 17W, Sec. 33	SH	RM-7.0	Shapavalov	Field note: "According to Warden Ovid Holmes and Mr. W. S. Harman of Fort Bragg, only straggler king salmon enter and spawn in both Big River and the Noyo river. Somewhat more enter and spawn in Ten Mile River but, according to the information provided by the above men, do not form a real run."	1948	
	COH	RM-7.0				
	CHIN	RM-7.0				
				Cherr & Griffin	Stream inventory: COH and SH were present in this stream.	1979
				Jones	Personal observation: COH and SH pass through this area to spawn in the headwater and tributaries and for the juveniles to migrate to sea. Have seen spring run CHIN, CDFG planted fish, holding in this area during the summer in the mid 1980's. The juvenile SH also utilize this area for rearing purposes. To my knowledge, this area was not used by juvenile COH for rearing.	1979/80
				Baker	Memo: 400,000 spring run CHIN were planted in the Ten Mile River. The source of the fish was the Trinity River Hatchery.	1980
				Welch	Memo: Spring run chinook salmon (introduced by CDFG) reported migrate through this area to spawn in the Middle Fork.	1982
			Salmon Restoration Association	Survey: Fall Run Chinook, 40 were caught in traps in the upper part of this section from October 8th, through December 8 th .	1985	
			Brown and Moyle	Report: COH reported in this stream; "Status of Coho Salmon in California", 1991	1991	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³	
South Fork Ten Mile River T19N, R17W, Sec. 3	COH	RM-14.0	Nye and Jones	Survey: found juvenile COH (30% of the total number of fish present in the lower stream area and 20% of the fish present in the headwater). A log jam limited the number of COH present in the drainage above Redwood Creek. This survey also found juvenile SH present (70% of fish present in the lower area and 80% in the headwater upstream of the log jam above Redwood Creek).	1961	
	SH	RM-17.5				
	CHIN	RM-??				
				Wilson and Steffens	Survey: found juvenile COH and SH near the bridge crossing on Camp Tenmile Road in early July.	1982
				Rhodes and Heckel	Survey: found juvenile COH and SH present between Campbell Creek and Churchman Creek. Between Churchman Creek and Redwood Creek juvenile COH and SH numbered 20-30 per mile. Above Redwood Creek only juvenile SH were present. They numbered 100/100 feet for the first 3 miles and then numbered 30/100 feet of stream up to a barrier. None were found above.	1982
				Jones	Survey: found juvenile COH and SH present. Electrofished: found only juvenile SH.	1983
				Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Maahs	Personal observation: 72 COH spawned above the instream trap. Juvenile COH were recorded migrating D/S through the migrant trap.	1995-96	
			Jones	Personal observation: CHIN have been observed in the South Fork Ten Mile River below Churchman Creek	1997	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Smith Creek T19, R17W, Sec. 11	COH	RM-3.5	Jones	Personal observations: juvenile COH and SH were present.	1979
	SH	RM-5.0	Ambrosius and Swirhun	Survey: "A few salmonids were observed"..., but the species were not identified.	1980
			Ubry and Plum	Survey: juvenile COH were present July 15 in numbers of 200/100 feet of stream.	1982
			Jones	Electrofished: found only juvenile SH above the sheep pasture.	1991
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
Maahs		Carcass survey: adult fish were counted during carcass surveys and juvenile COH and SH were trapped in a D/S migrant trap.	1995-96 1997		
Campbell Creek T19N, R14W, Sec. 11	COH	RM-3.5	Jones	Personal observations: juvenile COH and SH were present.	1979
	SH	RM-3.5	Davis and Sibbald	Survey: juvenile COH and SH were found in sparse to abundant numbers.	1982
			Jones	Electrofished: found only juvenile SH.	1991
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
Maahs		Carcass survey: COH found during carcass surveys. Downstream migrant surveys: juvenile COH and SH were trapped.	1995-96 1997		
Churchman Creek T19N, R16W, Sec. 20	COH	RM-2.0	Jones	Personal observations: both juvenile COH and SH were present.	1979
	SH	RM-2.0	Ambrosius and Swirhun	Survey: "salmonids were present", the species was undetermined.	1980
			Wilson and Steffens	Survey: found juvenile COH and SH present. In total, the salmonids numbered about 100/100 feet of stream.	1982
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
Maahs		Carcass survey: adult COH were observed.	1995-96		

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Unnamed Tributary to Churchman Creek T19N, R16W, Sec. 28	SH COH	RM-? RM?	Wilson and Steffens	Survey: found "salmonids", none were seen above jam # 4.	1982
Redwood Creek T19N, R16W, Sec. 23	COH SH	RM-2.3 RM-3.8	Crowdus & Santos Plumb and Ubry Sibbald and Davis Brown and Moyle	Survey: juvenile COH were present (35% of the salmonids below the middle fork). The survey also found juvenile SH present (65% of the salmonids present below the Middle Fork). Above the Middle Fork only juvenile SH were present. Survey: found juvenile COH present in Redwood Creek. Survey: found juvenile SH and COH present in the headwaters, above the unnamed tributary in section 12 in numbers of 25/100 feet of stream. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1961 1982 1982 1991
North Fork Redwood Creek T19N, R16W, Sec. 14	SH	RM-1.5	Crowdus and Santos Ubry and Plumb	Survey: juvenile SH were present. Survey: no fish were seen.	1961 1982
Unnamed Tributary to Redwood Creek (Middle Fork) T19N, R16W, Sec. 12	COH SH	RM-? RM-1.0	Crowdus Santos Sibbald and Davis	Survey: juvenile SH were present. Survey: juvenile COH and SH were present. The COH population did not extend above the mouth of the stream.	1961 1982
Mill Creek T20N, R17W, Sec. 35	SH	RM-2.0	Jones Swirhun, Nicol and Litwin Aasen Grass	Personal observation: juvenile SH were present. Survey: juvenile SH present. The stream could use more pools. Electroshocked: found juvenile SH present.	1979 1980 1983

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Middle Fork Ten Mile River T20N, R17W, Sec.25	COH	RM-11.25	Jones	Personal observation: both juvenile COH and SH were present.	1979
	SH	RM-13.25	Baker	Memo: A total of 400,000 juvenile spring run salmon were planted by CDFG in the Ten Mile River.	1980
	CHIN	RM-??	Welch	Note to CDFG files: 21 redds were observed in the stream after the first fall rains. These redds were said to be spring run chinook redds.	1982
			Heckel and Wilson	Survey: found juvenile COH and SH present between the mouth of Bear Haven and Little Bear Haven Creeks.	1982
			Uby and Sibbald	Survey: juvenile COH and SH present between Booth Creek and the unnamed Tributary in Section 32. Salmonid populations were 250 fish/100 yards of stream.	1982
			Steffens and Rhodes	Survey: found juvenile COH and SH present in the headwater above the Unnamed Tributary in Section 32 in numbers of 120/100 feet in the lower reach and 1-2 in the upper reach.	1982
			Maahs	Field note: CHIN were trapped for spawning purposes Oct 8, to Dec 8 th .	1985
			SRA	Report: The Salmon Restoration Association caught a total of 36 (28 male and & 8 female) COH during upstream trapping.	1985
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Jones	Electrofished: found only juvenile SH present.	1991
			GP	Electrofished: juvenile SH were found in the upper portion of the stream.	1993
		Maahs	Carcass surveys: COH, CHIN and SH were observed.	1995/96	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Bear Haven Creek T20N, R16, Sec. 31	COH	RM-3.0	Jones	Personal observation: juvenile COH and SH were observed.	1979
	SH	RM-4.4	Swirhun	Survey: COH observed but scarce. Canopy and instream shelter was excellent.	1980
			GP	Electrofished: found only juvenile SH.	1983
			Jones	Electrofished: juvenile COH and SH were present.	1991
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			GP	Electrofished: no fish were observed.	1993
			GP	Electrofished: some juvenile COH were seen in the lower area.	1997
South Fork Bear Haven Creek T20N, R16W, Sec. 29	SH	RM-1.4	Jones	Personal observation: juvenile SH were present.	1979
Horsetail Gulch T20N, R16W, Sec. 4	?	RM-?	Rhodes and Steffens	Survey: no fish were seen.	1982

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Little Bear Haven Creek T20N, R16W, Sec. 33	COH	RM-2.7	Rowell & Adams	Survey: juvenile salmonids were present up to the barrier falls, some 10 feet in height 2.7 miles up.	1961
	SH	RM-2.7+	Swirhun and Nichol	Survey: no fish were observed. Two log jam barriers, one 100 feet above the mouth and the other 0.3 miles up.	1980
			Aasen & Grass	Electrofished: some juvenile SH were found.	1983
			Jones	Electrofished: found only juvenile SH.	1983
			Jones	Electrofished: found only juvenile SH.	1991
			Maahs	Carcass survey: no COH were observed. However some SH were found.	1991-92
			Maahs	Carcass survey: COH and SH were observed during the surveys.	1995-96
Booth Gulch T19N, R16W, Sec. 1	SH	RM-0.8	Jones	Personal observation: juvenile SH were present.	1979
			Jones and Kidd	Electrofished: found juvenile SH present in densities of 1.0 fish/m ²	1983
Unnamed Tributary to Middle Fork Ten Mile River T19N, R15W, Sec. 6	COH	RM-?	Davis and Plumb	Survey: found juvenile COH present in this stream, extent unknown.	1982
Unnamed Tributary to Middle Fork Ten Mile River	Brown trout	RM-?	Plumb and Wilson	Survey: found Brown trout (resident) in abundant numbers up to jam# 6. This population was above a 60 foot falls.	1982

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
North Fork Ten Mile River T20N, R17W, Sec. 25	COH	RM-11.75	Jones	Personal observations: found juvenile COH and SH present.	1979
	SH	RM-14.5	Steffens and Wilson	Survey: both juvenile COH and SH were present. The juvenile COH numbered 10 / 100 feet of stream and the juvenile SH numbered 100/100 feet of stream.	1982
	CHIN	RM-?	Jones	Carcass survey: found three CHIN.	1989/90
			Jones	Electrofished: found only juvenile SH.	1991
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Maahs	Electrofished: COH pass upstream to the Little North Fork and Bald Hill Creek tributaries. Electrofishing revealed the juvenile SH density was 0.17fish/m ²	1993
			Jones	Personal observations: CHIN have been observed in this stream.	1982
			Jones and Sprig	Carcass survey: found three CHIN.	1988/89
Little North Fork Ten Mile River T20N, R17W, Sec. 14.	COH	RM-2.9	Rowell and Adams	Survey: large numbers of juvenile COH and SH were present.	1961
	SH	RM-3.5	Jones	Personal observation: found juvenile SH up to the foot of a dam.	1979
	CHIN	RM-?	Swirhun, Nicol and Litwin	Survey: juvenile SH were present and were common to abundant.	1980
			Jones	Electrofished: found only juvenile SH.	1983
			Maahs	Electrofished: the juvenile SH density was 0.80fish/m ² .	1992
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Maahs	Carcass survey: found COH and 1 CHIN present.	1991-92
			Maahs	Carcass survey found COH and SH present.	1995-96

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Buckhorn Creek T20N, R17W, Sec. 13	SH	RM-1.3	Rowell and Adams	Survey: found only juvenile SH. A large log jam at the mouth was suspected to have stopped COH from entering.	1961
	COH	RM-?	Davis and Wilson	Survey: found juvenile COH and SH present. Juvenile COH were "fairly plentiful at lower reaches".	1982
			Jones and Kidd	Electrofished: found only juvenile SH.	1983
			Jones	Electrofished: found only juvenile SH.	1984
			Jones	Electrofished: found only juvenile SH.	1991
Barlow Gulch T20N, R17W, Sec. 15	none	RM-0	Heckel and Sibbald	Survey: no fish were seen.	1982
Blair Gulch T20N, R17W, Sec. 11	none	RM-0	Plumb and Rhodes	Survey: no fish were seen.	1982
Unnamed Tributary to Little North Fork Ten Mile River T20N, R17W, Sec. 2	COH	RM-?	Sibbald and Heckel	Survey: found both juvenile COH and SH present.	1982
	SH	RM-?			
Cavanaugh Gulch T20N, R16W, Sec. 18	none	RM-0	Sibbald and Ubry	Survey: no fish were found.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Bald Hill Creek T20N, R16W, Sec. 9	COH	RM-2.0	Snyder	Memo: Removal of a barrier near the mouth opened 2.0 miles of stream to COH.	1980
	SH	RM-2.0	Yates and Swirhun	Survey: found juvenile SH present. A 40 foot bedrock waterfall was located about 100 yards below the road crossing on the East Fork of the Unnamed Tributary..	1980
			Kidd	Electrofished: found 13 juvenile SH in two passes, no COH.	1983
			Jones	Electrofished: found only juvenile SH.	1991
			Maahs	Electrofished: found juvenile COH and SH.	1993
Unnamed Tributary to North Fork Ten Mile River (Section 14 Creek) T20N, R16W, Sec. 14	COH	RM-?	Uby and Sibbald	Survey: juvenile COH were present, but the extent of the population was undetermined.	1982
Unnamed Tributary to North Fork Ten Mile River (Section 24 Creek) T20N, R16W, Sec. 24	COH SH	RM-? RM-?	Uby and Sibbald	Survey: both juvenile COH and SH were present, however the extent of the fishery use in the 1.1 miles of stream surveyed was not listed.	1982
Patsy Creek T20N, R15W, Sec. 18	SH	RM-0.7	Jones	Personal observation: juvenile SH were present.	1979
	COH	RM-?	Heckel	Survey: juvenile COH and SH were present. An abundant fingerling population was noted.	1982
			GP	Found juvenile SH, no COH.	1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Unnamed Tributary to North Fork Ten Mile River T20N, R15W, Sec. 17	COH SH	RM- 30 feet RM-0.1	Rhodes and Heckel	Survey: found both juvenile COH and SH above the mouth of this stream.	1982
Unnamed Tributary to North Fork Ten Mile River T20N, R15W, Sec. 17	COH	RM-?	Davis and Sibbald	Survey: found COH in numbers of 17/100 feet of stream near the mouth.	1982
Stanley Creek T20N, R15W, Sec. 17	?	RM-?		A stream survey is needed.	
Mill Creek T19N, R17W, Sec 19	SH COH	RM-2.5 RM-?	Crowdus & Santos Brown and Moyle Harris	Survey: the stream passes through Lake Cleone near it's confluence with the pacific Ocean. The lake level is controlled by a weir. Juvenile SH were found upstream of the lake. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: found juvenile SH present at a point upstream of the lake and downstream of the State HWY 1 bridge.	1961 1991 1997
Virgin Creek T19N, R17W, Sec. 31	SH	RM-1.6	Jones	Survey: juvenile SH were present, no COH. Tidewater goby were also present in the lagoon.	1961

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Pudding Creek T18N, R18W, Sec.1	COH	RM-10.0	Jones	Personal observation: juvenile COH and SH were present. CDFG had a Pudding Creek COH egg taking station in operation from the late 1950's until early 1960's. Following this, the egg taking operation was moved to the South Fork Noyo River. Survey: both juvenile COH and SH were present.	1959
	SH	RM-12.5	Primbs		1963
			Jones	Electrofished: 1983-to present. Standing crop density and biomass information is available through CDFG.	1983- 1997
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Harris	Electrofished: juvenile COH and SH were present. Carcass survey: 10 miles, found COH carcasses and 4 redds.	1998 1999/00
Noyo River T18N, R18W, Sec 13	COH	RM-30.0	Galigher Elwell	Survey: juvenile COH and SH were present.	1957
	SH	RM-30.0	Shapovolov	Field note: "According to Warden Ovid Holmes and Mr. W.S. Harman of Fort Bragg, only straggler king salmon enter and spawn in both Big River and Noyo river. Somewhat more enter and spawn in Ten Mile River but, according to the information provided by the above men, do not form a real run".	1948
	CHIN	RM-?			
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
		Jones	Personal observations: CHIN have been trapped in the CDFG station on the South Fork from time to time.	1997	
Hayshed Gulch T18N, 17W, Sec. 15.	COH	RM-0.5	Cherr Griffin	Fishery inventory indicates both COH and SH were present. A stream survey is needed.	1979
	SH	RM-0.5			

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
South Fork Noyo River T18N, R17W, Sec. 14	COH	RM-8.4	Jones	Personal observation: COH and SH use this drainage for spawning and rearing purposes. A COH trapping station has been built and operated on this stream since the early 1960's. In addition downstream migrant trapping is conducted each spring to assess populations of juvenile COH and SH. One CHIN was found as a carcass during the 1987-88 operation of the coho salmon egg taking station and three were caught during the 1997-98 operational season. When caught at the Station the fish were passed over the dam to proceed upstream and spawn. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Personal observation: A CHIN carcass was observed during a carcass survey below the Station on February 11, 2000. Carcass survey: three surveys of the mainstem SF (Dec, Jan & Mar.) Counted 29 live COH and 18 redds. Downstream migrant trapping (3 years) found juvenile COH and SH each year.	1997
	SH	RM-8.4	Grass		1987-88 1997-98
	CHIN	RM-3.5+	Brown and Moyle		1991
			Jones		2000
			Harris		1999-2000
			Harris		1998, 1999, 2000
Kass Creek T18N, 17W, Sec. 14	COH	RM-2.5	Primbs	Survey: juvenile COH were netted, no SH were seen.	1967
	SH	RM-2.5+	Litwin and Swirhun	Survey: juvenile COH and SH were observed.	1981
			Moore	Field note: in the CDFG response to timber harvest plan 1-86-16, COH and SH were noted present in the stream.	1986
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Jones	Personal observations: COH and SH were present.	1999

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
North Fork South Fork Noyo River T18N, R16W, Sec. 30	COH	RM-6.0	Fender, Litwin and Swirun	Survey: juvenile COH and SH were “common” in lengths of 2 to 3 inches.	1980
	SH	RM-?	Brown and Moyle	Report: COH were reported in this stream; “Status of Coho Salmon in California”, 1991	1991
			Jones	Personal observation: Both COH and SH were present this year in this drainage. In recent years a downstream trapping station has been set up to enumerate the seaward migration of juvenile COH and SH. The data is available from CDFG.	1997
Brandon Gulch T18N, R16W, Sec. 19	SH	RM-0.4	Mesman & Dombrowski	Electrofished: 2 sites, found juvenile SH 2,237 feet upstream.	1996
Little North Fork Noyo River T18N, R16W Sec. 6	COH	RM-3.5	Jones	Personal observation: both COH and SH were present and use this drainage.	1961-1997
	SH	RM-3.5	Brown and Moyle	Report: COH were reported in this stream; “Status of Coho Salmon in California”, 1991	1991
Bear Gulch T18N, R16W, Sec. 32	COH	RM-0.7	Hichethier & Mesman	Electrofished: 5 sites, found juvenile COH and SH 3,772 feet upstream.	1995
	SH	RM-0.7			
Parlin Creek T17N, R16W, Sec. 4	COH	RM-3.0	Morehouse	Survey: juvenile COH and SH were present in the lower 3 miles.	1959
	SH	RM-3.6	Brown and Moyle	Report: COH were reported in this stream; “Status of Coho Salmon in California”, 1991	1991
			Coyle & Young	Electrofished: 5 sites, juvenile COH were present up to 15,856 feet. Juvenile SH were present to 18,797 feet upstream.	1995
Duffy Gulch T18N, R16W, Sec.12	SH	RM-1.4	Jones	Personal observation: SH have been seen in this stream.	1961
	COH	RM-1.7	Fender, Litwin and Swirhun	Survey: found only COH but the population was said to be “scarce”.	1980
			Brown and Moyle	Report: COH were reported in this stream; “Status of Coho Salmon in California”, 1991	1991

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Gulch Thirty one T18N, R15W, Sec.7	COH SH	RM-0.6 RM-0.6	Jones	Personal observation: both COH and SH have been seen in this stream.	1961
Unnamed Tributary # 6 T18N, R15, Sec.17	COH SH	RM-1.0 RM-1.0	Jones Daugherty	Personal observation: juvenile COH and SH were seen in this stream. Electrofished: found juvenile SH, no COH	1961 1996
North Fork Noyo River T18N, R15W, Sec.8	COH SH	RM-5.7 RM-5.7	Jones Brown and Moyle	Personal observation: Juvenile COH and SH were present in this stream in good numbers. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1961 and 1997 1991
Marble Gulch T18N, R15W, Sec 9	COH SH	RM-1.0 RM-1.0	Jones Fender and Litwin Brown and Moyle Gilroy & Beaumont	Personal observation: juvenile COH and SH were seen in this stream. Survey: juvenile COH were present in the lower 0.2 miles. No other fish were seen. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofishing: 6 sites, juvenile SH were present up to 1,155 feet.	1961 1980 1991 1999
Gulch Seven T18N, R115W, Sec.3	COH SH	RM-0.5 RM-0.5	Jones	Personal observation: juvenile COH and SH were seen in this drainage.	1961
Hayworth Creek T19N, R15W, Sec. 33	COH SH	RM-3.0 RM-3.0	Primbs Fender and Litwin Brown and Moyle Gilroy, Taylor & Jankowski	Survey: juvenile COH and SH were present. Survey: found juvenile SH present. The density of these fish was 10/ 100 feet of stream in the area from the mouth to the North Fork and 20/100 feet of stream from the North Fork to Soda Creek. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: 13 sites, juvenile SH were found up stream 23,765 feet.	1967 1980 1991 1999

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
North Fork Hayworth Creek T19N, R15W, Sec. 26	COH	RM-1.3	Gallagher	Survey: juvenile COH and SH were present.	1957
	SH	RM-1.3	Jones	Personal observation: found juvenile SH, "small fish 1 to 3 inches but not abundant".	1961
			Swirhun and Litwin	Survey: found COH present below the 10 foot high rock and log barrier located some 300 yards above the mouth.	1980
			Daugherty	Electrofished: juvenile SH were present.	1996
			Gilroy & Beaumont	Electrofished: 4 sites, found juvenile SH present up stream 4,383 feet.	1999
Soda Creek T19N, R15W, Sec. 25	SH	RM-0.1	Gilroy, Turner & Jankowski	Electrofished: 8 sites. Juvenile SH were found upstream 304 feet.	1999
Panther Creek T19N, R15W, Sec. 24	SH	RM-0.3	Gilroy & Beaumont	Electrofished: 6 sites, found juvenile SH present 1,389 feet upstream.	1999
Middle Fork of North Fork T19N, R15W, Sec. 29	SH	RM-2.0	Galligher	Survey: juvenile SH were present, found a barrier on the North Fork downstream of the Middle Fork.	1957
	COH	RM-?	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Daugherty	Electrofishing: found juvenile SH.	1994, 95, 96
			Gilroy, Turner, & Jankowski	Electrofished: 15 sites, juvenile SH were present up stream 1,596 feet.	1999
Dewarren Creek T19N, R15W, Sec. 29	SH	RM-2.0	Galligher	Survey: found a barrier on the North Fork downstream of the Middle Fork. Juvenile SH were present below.	1961
			Daugherty	Electrofished: juvenile SH were present high in the drainage.	1994, 1995, 1996
			Gilroy & Beaumont	Electrofished: 8 stations, SH were found up to 549 feet upstream.	1999

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Olds Creek T18N, R17 Sec. 6	COH	RM-5.6	Jones	Personal observation: juvenile COH and SH were present.	1961
	SH	RM-6.0	Brown and Moyle Daugherty	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: only juvenile SH were present.	1991 1996
Unnamed Tributary # 7 T18N, R15W, Sec. 15	COH	RM-0.8	Jones	Personal observation: juvenile COH and SH were present.	1961
	SH	RM-0.8	Daugherty	Electrofished: juvenile SH were present (just above the mouth).	1996
Redwood Creek T18N, R15W, Sec. 11	COH	RM-3.5	Jones	Personal observation: juvenile COH and SH were present.	1961
	SH	RM-4.0	Brown and Moyle Daugherty	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: COH and SH were present.	1991 1996
Gulch C T18N, R15W, Sec. 12	SH	RM-1.0	Jones	Personal observation: juvenile SH were present.	1961
McMullen Creek T18N, R14W, Sec. 7	SH	RM-0.7	Jones Daugherty	Personal observation: juvenile SH were present. Electrofished: juvenile SH were present both years.	1961 1995, 1996
Burbeck Creek T18N, R15W, Sec. 8	SH	RM-1.0	Jones Daugherty	Personal observation: juvenile SH were present. Electrofished: juvenile SH were found.	1961 1996
Hare Creek T18N, R18W, Sec. 13	COH	RM-9.6	Swirhun	Survey: salmonids were present in low numbers but were not identified to species.	1981
	SH	RM-8.4	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Mesman & Hichethier	Electrofished: 7 sites, juvenile SH were present 50,566 feet upstream. COH were present 44, 460 feet upstream.	1995
			Harris	Electrofished: both COH and SH were present 8 miles upstream.	1995-96-97

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Covington Gulch T18N, R17W, Sec. 20	SH	RM-.01	Harris	Electrofished: juvenile SH were present.	1996
Walton Gulch T18N, R17W, Sec. 27	COH SH	RM-0.1 RM-0.05	Jones Dunn & Bettina	Electrofished: COH and SH were present. The fishway was in poor condition. Electrofished: juvenile SH were present 270 feet upstream.	1988 1995
South Fork Hare Creek T18N, R17W, Sec. 27	COH SH	RM-0.5 RM-0.7	Brown and Moyle Dunn & Chimarios Harris	Report: COH reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished 6 sites: juvenile SH were present 2,827 feet upstream. Electrofished: juvenile SH were present 0.3 miles upstream.	1991 1995 1996
Bunker Gulch T18N, R17W, Sec. 26	COH SH-	RM-1.0 RM-1.6	Swirhun Brown and Moyle Harris	Survey: No fish were observed, but "have been seen by the local CDFG warden". Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: juvenile COH and SH were present.	1981 1991 1996
Digger Creek T18N, R18W, Sec. 24	SH	RM-0.5	Jones	Personal observation: the culvert at the Hwy 1 crossing was a barrier.	1967
Mitchell Creek T18N, R18W, Sec. 25	COH SH	RM-0.3 RM-2.8	Jones Wdn. Bain	Personal observation: observed marked hatchery coho, Pudding Creek origin, upstream of the HWY 1 bridge. Memo: a timber harvest violation report by CDFG Warden Bain occurred on this stream.	1961 1973
Jug Handle Creek T18N, R18W, Sec. 36	SH COH	RM-4.0 RM-?	Brown and Moyle Harris	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: Juvenile SH were present. However, COH were once present some years ago (Jones).	1991 1995-96-97

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Casper Creek T18N, R17N, R17W, Sec. 1	COH	RM-3.8	Harris	Electrofished: carcass surveys, electrofishing and downstream migrant trapping data is available from CDFG. Juvenile and adult COH and SH were present in this drainage.	1983-2000
	SH	RM-3.8	Mesman & Hichethier	Survey: COH & SH were found up to 18, 849 feet above the mouth.	1995
			Colins	Survey: found 21 juvenile COH and 96 juvenile SH at a site 13,887 feet upstream and 20 juvenile COH and 104 juvenile SH at 14,793 feet upstream.	1999
South Fork Caspar Creek T17N, R17W, Sec. 9	COH	RM-1.6	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
	SH	RM-2.3	Mesman & Hichethier	Electrofished: 4 sites, found juvenile COH and juvenile SH at 3,096 feet upstream and juvenile SH again at 12,595 feet upstream.	1995
			Harris	Electrofished: juvenile COH and SH were present.	1996
North Fork Caspar Creek T17N, R17W, Sec. 9	COH	RM-2.7	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
	SH	RM-3.9	Harris	Electrofished: Juvenile COH and SH were present. Carcass data also revealed these species present.	1966
Unnamed tributary to North Fork Caspar Creek (Middle Fork) T17N, R17W, Sec. 10	COH	RM-0.7	Mesman, Hichethier & Chimarios	Electrofished: found juvenile SH up to 2,389 feet, no COH.	1995
	SH	RM-1.0	Harris	Electrofished: found juvenile COH and SH present.	1996

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Doyle Creek T17N, R17W, Sec. 1	COH	RM-0.2	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
	SH	RM-1.0	Jones	Personal observation: including electrofishing, both juvenile COH and SH were found below the Hwy 1 crossing. The HWY 1 culvert was a barrier to COH, none were found above. As there is no spawning habitat in the lower stream reach I suspect the juvenile COH present, enter the stream from Caspar Creek by way of the ocean.	1996
			Daugherty	Electrofished: found juvenile SH or possibly resident RT present upstream of the Cal Trans culvert.	1995, 1996
			Harris	Electrofished: found juvenile SH, but no COH in October.	1999
Russian Gulch T17N, R17W, Sec. 18	COH	RM-2.2	Jones	Personal observation: including electrofishing, found both juvenile COH and SH present.	1983-1997
	SH	RM-2.2	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Harris	Carcass survey: Found SH only, (mouth to waterfall).	1998/99
South Fork Russian Gulch T17N, R17W, Sec. 20	COH	RM-0.6	Jones	Personal observation: juvenile COH and SH have been seen in this drainage.	1983
	SH	RM-0.6			
Jack Peters Gulch T17N, R17W, Sec. 19	SH	RM-0.9	Week	Survey: juvenile SH were present in the lower stream. The diversion dam owned by Surfwood Estates was believed to be a barrier. No fish were seen above.	1974

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Big River, T17N, R17W, Sec. 30	COH	RM-39.6	Shapovolov	Field note: "according to Warden Ovid Holmes and Mr. W.S. Harman of Fort Bragg, only straggler king salmon enter and spawn in both Big River and Noyo River. Somewhat more enter and spawn in Ten Mile River but, according to the information provided by the above men, do not form a real run".	1948
	SH	RM-41.3			
	CHIN	RM ?			
				U.S. Fish and Wildlife Jones Brown and Moyle Daugherty	Survey: Fisheries Improvement Study, Big River (Eureka division, North Coast Project, California), electrofishing confirmed the presence of both COH and SH. Personal observation: found COH, SH and CHIN present in the mid 1980's. Found juvenile COH and SH throughout the major portion of the tributaries in the lower drainage during the years of 1958 through 1997. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: only juvenile SH were present at a site below Valentine Creek, but juvenile COH were found just upstream of the confluence of Russell Brook Creek in 1996.
Railroad Gulch T17N, R17W, Sec. 24	COH	RM-1.3	Harris	Electrofished: found juvenile COH present, no SH in 1995. However, juvenile SH were present in 1996 and 1997.	1995-96-97
Little North Fork Big River T17N, R17W, Sec. 24	COH	RM-6.6	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
	SH	RM-7.0	Mesman,& Hickethier	Electrofished: 3 sites, found juvenile COH up to 10,603 feet and juvenile SH up to 19,341 feet.	1995
			Harris	Electrofished: juvenile COH and SH were present.	1995-96-97
			Harris	Carcass surveys: found redds but no carcasses or live fish both years.	1998/99 &1999/00
Thompson Gulch T17N, R16W, Sec. 18	COH	RM-0.5	Harris	Electrofished: juvenile COH and SH were present in 1995. No SH were present in 1996 but were again present in 1997.	1995-96-97
	SH	RM-0.1	CCC	Electrofished: 3 sites, juvenile COH up to 3,442 feet with no juvenile SH at that site. Juvenile SH were present in the lower area up to 316 feet.	1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
East Branch Little North Fork Big River T17N, R16W, Sec. 8	COH	RM-1.8	Jones	Personal observation: juvenile COH and SH were present.	1967
	SH	RM-1.8	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
Berry Gulch T17N, R16W, Sec. 8	COH	RM-0.8	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
	SH	RM-1.4	Harris	Electrofished: found juvenile COH and SH up to 0.8 miles.	1995-96-97
			Cooper & Mesman	Electrofished: 4 sites, juvenile COH were present up to 436 feet, juvenile SH extended up to 7,164 feet.	1997
Two Log Creek T17N, R16W, Sec. 23	COH	RM-3.3	Thomas	Survey: only juvenile SH were present.	1966
	SH	RM-3.3	Jones	Electrofished: COH and SH were present.	1983
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			MacMillan and Retherford	Electrofished: 4 sites, juvenile COH and SH were found up to 17,658 feet.	1998
			Daugherty	Electrofished: found both juvenile COH and SH present.	1995
		Harris	Electrofished: found juvenile COH and SH present. Snorkeled: juvenile COH and SH were present in July.	1996 1995-96-97 2000	
Tramway Gulch T17N, R16W, Sec. 24	SH	RM-0.6	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Daugherty	Electrofished: found juvenile SH in mid Sec. 24 both years. A bank failure (earth slide) was present upstream.	1995 1996

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
North Fork Big River T17N, R15W, Sec. 29	COH	RM-8.8	Harris	Electrofished: both juvenile COH and SH were present.	1966
	SH	RM-13.2	U.S. Fish and Wildlife Jones	Electrofished: found juvenile COH and SH present.	1974
				Personal observation: both juvenile COH and SH were present.	1967
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Mesman & Cooper	Electrofished: 2 sites, found juvenile SH up to 54,167 feet (10.3 miles).	1997
			Daugherty Harris	Electrofished: found both juvenile COH and SH upstream of Harris Gulch in Sec. 17. Electrofished: found juvenile COH and SH present all three years.	1996 1995-96-97
East Branch North Fork Big River T17N, R15W, Sec. 20	COH	RM-5.0	Edie	Survey: Both COH and SH were present.	1966
	SH	RM-5.3	Brown and Moyle Harris	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: found juvenile COH absent and SH present.	1991 1995-96-97
			Retherford, Behm & Mac Millan	Electrofished: 5 sites, found juvenile SH up to 28,072 feet.	1998
			Daugherty	Electrofished: found both juvenile COH and SH present in section 15.	1996
Bull Team Gulch T17N, R15W, Sec. 15	COH	RM-0.2+	Daugherty	Electrofished: found both juvenile COH and SH present. Unsure of the length of habitat available to these fish above the electrofishing site.	1996
	SH	RM-0.2+			

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Chamberlain Creek T17N, R15W, Sec. 5	COH	RM-3.5	Jones and Yates	Survey: found juvenile COH and SH present. The COH numbered 30/ 100 feet of stream and the SH numbered as high as 200/100 feet of stream but averaged about 50 / 100 feet of stream.	1980
	SH	RM-4.6	Jones	Electrofished: both juvenile COH and SH were present.	1983
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Harris & Jones	Electrofished: Both juvenile COH and SH were present. The pinched bedrock area under the road crossing was a barrier to COH during low flow years.	1995-96-97
			Cooper & Mesman	Electrofished: 2 sites, found juvenile SH up 16,018 feet.	1997
Water Gulch T17N, R15W, Sec. 5	COH	RM-1.4	Montoya and Litwin	Survey: found "COH/SH" in this stream. The population in the lower reach was 40 fish/100 feet of stream.	1981
	SH	RM-2.0	Harris and Jones	Electrofished: juvenile COH and SH were present all three years.	1995-96-97
			Harris and Jones	Electrofished: juvenile COH and SH were present at 3,075 feet. The juvenile SH population extended up to 7,434 feet.	1997
Park Gulch T17N, R15W, Sec. 5	SH	RM-0.7	Analasiou and Swirhun	Survey: no fish observed. Found a rock falls barrier at 0.5 miles.	1981
			Jones	Electrofished: found juvenile SH present.	1983
			Harris	Electrofished: juvenile SH were present in 1995. No fish were seen in 1996. Juvenile SH were again present in 1997.	1995-96-97
			Mesman & Cooper	Electrofished: 1 site, juvenile SH were found 28 feet above the mouth.	1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
West Fork Chamberlain Creek T17N, R15W, Sec. 5	COH	RM-1.1	Anastasiou and Swirhun	Survey: none observed. Six barriers and 4 log jams were found.	1981
	SH	RM-3.0	Harris & Jones	Electrofished: juvenile COH and SH were present all three years.	1995-96-97
			Mesman & Cooper	Electrofished: juvenile SH were found 7,527 feet upstream.	1997
Gulch Sixteen T18N, R15W, Sec. 31	SH	RM-0.5	Harris	Electrofished: juvenile SH were present upstream 0.5 miles.	1995-96-97
			Cooper & Mesman	Electrofished: 2 sites, juvenile SH were found up to 1,606 feet.	1997
Arvola Gulch T18N, R15W, Sec. 28	COH	RM-0.5	Jones and Yates	Survey: juvenile COH and SH were present. Also found "15 log jams of which more than half were barriers".	1980
	SH	RM-1.0	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Harris & Jones	Electrofished: juvenile COH and SH were present all three years.	1995-96-97
Unnamed Tributary to Chamberlain Creek T18N, R15W, Sec. 28	SH	RM-0.4	Jones and Yates	Survey: a few juvenile SH were found in the lower end of the stream. There were 35 log jams and 10 barriers seen during the survey.	1980
			Harris & Jones	Electrofished: found juvenile SH present each year.	1995-96-97

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
James Creek T17N, R15W, Sec. 11	COH SH	RM-0.5 RM-4.0	Yates and Swirhun Brown and Moyle Harris Mesman & Wrights	Survey: juvenile "SH ranging from 2 to 5 inches were abundant throughout the stream, COH were reported to be present in the winter and spring". Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: juvenile COH and SH were present in 1996. COH were stopped by a barrier. Juvenile SH were also present in 1995 and 1997. Electrofished: 3 sites, only juvenile SH were present and found upstream to 19,975 feet.	1980 1991 1995-96-97 1996
North Fork James Creek T18N, R15W, Sec. 35	SH COH	RM-2.3 RM-?	Brown and Moyle Cooper & Mesman Harris	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: juvenile SH were found up to 9,585 feet upstream. Electrofished: juvenile SH were present for 2.3 miles.	1991 1995 1995-96-97
South Fork Big River T17N, R15W, Sec. 33	COH SH	RM-17.5 RM-?	Edie Brown and Moyle Mesman & Patton Jones Daugherty	Survey: juvenile COH and SH were present. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: 3 sites, found juvenile SH up to 32,856 feet . Electrofished: Juvenile SH were present but COH were not seen above Daugherty Creek. Electrofished: juvenile COH were found in Sec. 34, of T17N, R15W, and were known to use the tributaries above.	1966 1991 1995 1996 1996

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Ramon Creek T16N, R15W, Sec. 2	SH COH	RM-3.7 RM-1.2	Cherr & Griffin Brown and Moyle Daugherty	Stream inventory : data indicates SH were present. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: found juvenile COH and SH present. The COH were present into Sec. 1, 200 feet above the confluence of the North Fork of Ramon Creek. The juvenile SH were present above.	1979 1991 1995
North Fork Ramon Creek T16N, R15W, Sec. 1	COH SH	RM-0.5 RM-0.5	Daugherty	Electrofished: juvenile COH and SH were present.	1995
Mettick Creek T16N, R15W, Sec. 10	SH	RM-0.7	Daugherty	Electrofished: juvenile SH were present all three years.	1994, 1995, 1996
Anderson Gulch T16N, R15W, Sec. 11	SH	RM-200 feet	Daugherty	Electrofished: found juvenile SH present all three years.	1994, 1995, 1996
Half Way House Gulch T16N, R15W, Sec. 14	SH	RM- 500 feet	Daugherty	Electrofished: juvenile SH were present.	1996
Daugherty Creek T16N, R14W, Sec. 19	COH SH	RM-2.7 RM-7.0	Brown and Moyle Mesman & Patton Daugherty	Report: COH reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: 3 sites, found only juvenile SH present, up to 32,856 feet (6.2 miles). Electrofished: found juvenile COH 2.7 miles and SH 7.0 miles	1991 1993 1996
Soda Creek T16N, R14W, Sec. 29	SH	RM-1.2	Daugherty Cooper & Mesman	Electrofished: found SH present all three years. Electrofished: 3 sites, no COH or SH were present.	1995, 1996 1997 1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Gates Creek T16N, R14W, Sec 32	SH	RM-2.5	Mesman and Patton Daugherty	Electrofished: 3 sites, juvenile SH were present up to 10,958 feet. Electrofished: juvenile SH were present at 2.5 miles.	1993 1996
Johnson Creek T16N, R14W, Sec. 33	SH COH	RM-1.2 RM-?	Brown and Moyle Daugherty	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: found juvenile SH present.	1991 1996
Snuffins Creek T15N, R14W, Sec. 10	SH	RM-1.2	Mesman & Patton Daugherty	Electrofished: 3 sites, juvenile SH were present up to 860 feet. Electrofished: found juvenile SH present to 1.2 miles.	1993 1996
Johnson Creek T16N, R14W, Sec. 20	SH	RM-1.4	Holman, Rowell, Day, Morehouse and Nye Jones	Survey: found juvenile SH. The culvert problem at the mouth was believed to prevent COH from entering this stream. Personal observation: the problem, a three foot jump into a flat concrete box culvert, still exists in 2000. No COH currently use this stream although Nat Bingham reared and released young of the year COH to yearling size in this stream for many years.	1959 2000
Dark Gulch T16N, R14W, Sec. 16	SH	RM-0	Elwell Jones	Survey: no fish were seen. Personal observations: The county road culvert was a barrier, it is 80+ feet long, is paved and is at a grade of more than 2%. Stream flow characteristics do not fit the "Alaskan curve" for fish passage. Baffles are needed in the barrel of this culvert.	1958 1999
Montgomery Creek T16N, R14W, Sec. 22	SH	RM-0.4	Jones	Personal observation: A barrier falls currently blocks fish passage in the Montgomery Woods area. No modification is recommended as the gradient is steep above the falls.	2000

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Russell Brook Creek T17N, R15W, Sec. 26	SH	RM-3.5	Cherr & Griffin Jones Daugherty	Inventory: SH were present. Personal observation: juvenile SH were present. Electrofished: juvenile SH were present in the east side of Section 31.	1979 1967 1996
Pigpen Gulch T17N, R14W, Sec. 19	SH	RM-1.0	Cherr & Griffin Jones Daugherty	Inventory: SH were present. Personal observation: juvenile SH were seen. Electrofished: found juvenile SH above the mouth.	1979 1967 1994, 1995, 1996
Martin Creek T17N, R14W, Sec. 20	SH	RM-1.7	Cherr & Griffin Jones Daugherty	Inventory: SH were present. Personal observation: heavy logging damage was found in 1967. Electrofished: found juvenile SH present above the mouth all three years.	1979 1967 1994, 1995, 1996
Valentine Creek T17N, R14W, Sec. 33	SH	RM-2.4	Cherr & Griffin	Inventory: SH were present.	1979
Rice Creek T17N, R14W, Sec. 27	SH	RM-1.5	Nye & Morehouse Jones	Survey: no fish were seen. Personal observation: Juvenile SH were present. Another survey is needed.	1959 1967
Little River T16N, 17W, Sec. 6	COH SH	RM-4.5 RM-4.5	Brown and Moyle Primbs Jones Harris	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Survey: juvenile COH and SH were present. Carcass surveys, electrofishing and downstream migrant trapping: has been conducted by CDFG since 1983. Data on COH and SH findings are available from CDFG. Electrofished: carcass surveys, downstream traps. COH present in 2000 but in very low numbers. The stream needs complexity.	1991 1965 1997 1998-00

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Buckhorn Creek T16N, R17W, Sec. 8	SH	RM-2.0	Marco & Steitz	Survey: found juvenile SH up to the falls. The waterfall 1/4 mile up the North Fork is a barrier.	1976
	COH	RM-?	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
Dark Gulch T16N, 17W, Sec. 20	SH	RM-0.5	Jones	Electrofished: juvenile SH population density of 0.17 fish/m ² were found.	1990
Albion River T16N, R17W, Sec. 28	COH	RM-14.0	Santos & Jones	Survey: juvenile COH and SH were present.	1961
	SH	RM-16.0	Jones	Electrofished: found juvenile COH and SH present.	1983
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Daugherty	Electrofished: juvenile COH and SH were present to the confluence of the North Fork, also juvenile SH were found in the headwater in Section 9.	1996
			Harris	Electrofished: found juvenile COH and SH present near the Fire station.	1996
Railroad Gulch T16N, R17W, Sec. 23	SH	RM-1.7	Jones	Survey: occasionally SH enter this stream but it has been known to dry up during low flow years.	1961
	COH	RM-?	Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Daugherty	Electrofished: found no fish. However, found juvenile COH present in an unnamed tributary.	1996
Pleasant Valley Creek T16N, R17W, Sec. 23	COH	RM-0.2	Daugherty	Electrofished: found both juvenile COH and SH in this stream.	1995 & 1996
	SH	RM-0.3			
Unnamed tributary to Railroad Gulch T16N, R16W, Sec. 25	COH	RM-0.09	Daugherty	Electrofished: found juvenile COH and SH in this small tributary.	1996
	SH	RM-0.09.			

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Unnamed tributary to Railroad Gulch T16N, R16W, Sec. 25	COH SH	RM-0.04 RM-0.04	Daugherty	Electrofished: found juvenile COH and SH in this small tributary.	1996
Slaughter House Gulch T16N, R17W, Sec. 14	COH SH	RM-0.09 RM-0.09	Daugherty	Electrofished: found both juvenile COH and SH present.	1996
Duck Pond Creek T16N, R17W, Sec. 13	COH	RM-0.09	Daugherty	Electrofished: found only juvenile COH.	1996
South Fork Albion River T16N, R16W, Sec.17	COH SH	RM-5.0 RM-6.6	Nye & Adams Primbs Brown and Moyle Daugherty McKernan & Cambell	Survey: both juvenile COH and SH were present. Survey: found juvenile COH and SH present. Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991 Electrofished: juvenile COH and SH were both present above Larmer Gulch. Also found juvenile COH upstream in mid Section 23. Electrofished: 1 site, juvenile SH were present to 34,000 feet above the mouth.	1961 1966 1991 1992 1998
Little North Fork South Fork Albion River T16N, R16W, Sec. 21	COH SH	RM-0.2 RM-0.3	Daugherty CCC's/WSP	Electrofished: found juvenile COH and SH present Visual survey: juvenile SH were present at 1,628 feet.	1996 1998
Larmer Gulch T16N, R16W, Sec. 23	SH	RM-0.04	Daugherty	Electrofished: juvenile SH were found 200 feet above the mouth.	1996

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Bull Team Gulch T16N, R16W, Sec. 23	SH	RM-0.02	McKernan & Campbell	Electrofished: 1 site, juvenile SH were found 115 feet above the mouth.	1998
			Daugherty	Electrofished: found juvenile SH present in this stream.	1996
Winery Gulch T16N R16W, Sec. 14	SH	RM-0.25	Daugherty	Electrofished: found juvenile SH 1/4 mile above the mouth.	1996
North Fork Albion River T16N, R16W, Sec. 11	COH	RM-4.0	Jones & Adams	Survey: both juvenile COH and SH were present. Low stream flows (many intermittent areas) were believed to have limited the populations of both COH and SH.	1961
	SH	RM-?	Edie	Survey: found juvenile COH and SH present.	1966
			Brown and Moyle	Report: COH were reported in this stream; "Status of Coho Salmon in California", 1991	1991
			Daugherty	Electrofished: found juvenile COH and SH in the southern portion of Section 2.	1994- 95- 96
Kaison Gulch T16N, R16W, Sec. 7	SH	RM-0.9	Daugherty	Electrofished: juvenile SH were found.	1996
Railroad Gulch T16N, R16W, Sec. 10	COH	RM-0.4	Daugherty	Electrofished: juvenile COH and SH were present in this stream.	1996
	SH	RM-0.4			
Tom Bell Creek T16N, R16W, Sec. 4	COH	RM- 0.04	Daugherty	Electrofished: 2 sites, juvenile COH were both found in the lower station. However, juvenile SH were present in both stations.	1996
	SH	RM-1.0			
Marsh Creek T16N, R16W, Sec. 12	SH	RM-?	Cunningham	Survey: no salmonids were seen during this years drought.	1977
	COH	RM-?		Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1991

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Big Salmon Creek T16N, R17W, Sec. 28	COH	RM-8.3	Crowdus & Santos	Survey: found both juvenile COH and SH present.	1961
	SH		Jones	Electrofished: found juvenile COH and SH present in 1983.	1983
			Jones	Electrofished: standing crop and biomass data was collected for both COH and SH.	1986
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1991
Little Salmon Creek T16N, R17W, Sec. 28	COH	RM-0.9	Jones	Survey: juvenile COH and SH were seen. (Numerous log jams were also found).	1961
	SH	RM-0.9			
Hazel Creek T16N, R17W, Sec. ?	SH	RM-1.5	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1991
	COH	RM-1.5	Jones	Personal observation: COH and SH use 1.5 miles of this stream.	1997
Navarro River T15N, R17W, Sec. 5	COH	RM-28.3	Rowell & Keller	Survey: COH and SH used this drainage.	1962
	SH	RM-28.3	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1991
	STB	RM-2.0	Jones and Harris	Downstream migrant trapping: caught juvenile COH and SH both years.	1995 and 1996
			Ornbaun	Striped bass caught while angling.	1959
Marsh Gulch T15N, R17W, Sec. 11	SH	RM-0.6	D. Jones & MacMillan	Electrofished: 1 site, juvenile COH and SH were present upstream 259 feet.	1997
	COH	RM-0.04	Daugherty	Electrofished: 2 sites, juvenile COH and SH were present 300 feet above the mouth. Also juvenile SH were again found at the upper site 0.6 miles upstream.	1996

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Murry Gulch T15N, R17W, Sec. 12	COH	RM-0.09	D. Jones & MacMillan	Electrofished: 2 sites, juvenile SH were found 1,846 feet above the mouth.	1996
	SH	RM-0.3	Daugherty	Electrofished: found juvenile COH and SH 0.09 miles upstream.	1995
Barton Gulch T15N, R16N, Sec. 7	SH	RM-0.09	Daugherty	Electrofished: juvenile SH were present.	1994, 1995, 1996
Flume Gulch T15N, R16W, Sec. 18	SH	RM-1.1	D. Jones & MacMillan	Electrofished: 3 sites, found juvenile SH present upstream 3,422 feet upstream.	1996
			Daugherty	Electrofished: found juvenile SH present in Section 18.	1996
Ray Gulch T15N, R16W, Sec. 7	SH	RM-1.0	Coyle	Personal observation: no fish were seen in this class 2 stream.	1994
			Daugherty	Electrofished: found juvenile SH present in the upper portion of section 8.	1995, 1996
Mustard Gulch T15N, R16W, Sec. 16	SH	RM-0.09	Daugherty	Electrofished: found juvenile SH present 500 feet above mouth.	1995
North Fork Navarro River T15N, R16W, Sec. 16	SH	RM-5.0	Adams & Moore	Survey: no COH were seen, juvenile SH were present throughout the stream.	1962
	COH	RM-?	Jones	Personal observation: juvenile SH were seen throughout the stream. COH were only seen in the lower reaches. However, the extent of the COH use was undetermined.	1970-1997
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones and Harris	Downstream migrant trapping: both juvenile COH and SH were present near the scout camp, both years.	1995-96-97
Dead Horse Creek T15N, R16W, Sec. 15	COH	RM-0.09	Daugherty	Electrofished: found juvenile COH and SH in the site above HWY 128. In 1996 only juvenile SH were found.	1995, 1996
	SH	RM-0.09			

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Tramway Gulch T15N, R16W, Sec. 23	SH	RM-0.7	Daugherty	Electrofished: found juvenile SH present 0.7 mile above mouth.	1995
Coon Creek T16N, R16W, Sec. 14	SH	RM-0.25	Daugherty	Electrofished: found juvenile SH present 1/4 mile above the mouth.	1994 and 1996
Flynn Creek T15N, R15W, Sec. 13	COH	RM-3.8	Jones	Electrofished: found juvenile COH and SH present near HWY 128 and above.	1983
	SH	RM-3.8	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Mesman, Coyle, d. Jones, & MacMillan	Electrofished: 3 sites, juvenile COH and SH were present up to 15,625 feet above the mouth.	1995
			Daugherty	Electrofished: found juvenile COH and SH present in Section 36.	1995, 1996
			Jones and Harris	Electrofished: found both juvenile COH and SH present.	1997
Camp 16 Gulch T15N, R16W, Sec.	COH	RM-0.3	Daugherty	Electrofished: 2 sites, juvenile COH and SH were found up as far as 0.3 miles.	1996
	SH	RM-0.3			
Tank 4 Gulch T16N, R16W, Sec. 35	COH	RM- in mouth+	Daugherty	Electrofished: 1 site, found juvenile COH and SH in the mouth of Tank 4 Gulch.	1996
	SH	RM- in mouth+			
South Branch North Fork Navarro River T15N, R15W, Sec. 7	COH	RM-?	Jones	Personal observation: found juvenile COH as well as SH present in the lower reach, the extent of the COH population was unknown.	1975
	SH	RM-17.0	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Flosi, Coyle, Chimarios, Dunn and Young	Electrofished: found juvenile SH up to 89,578 feet above the Boy Scouts summer camp impoundment.	1995

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Bailey Creek T15N, 15W, Sec. 13	SH	RM-1.3	Coyle, Chimarios and Dunn	Electrofished: 3 sites, juvenile SH were found 7,024 above the mouth.	1995
Bear Creek T15N, R14W, Sec. 19	SH	RM-?	Flosi, Chimarios, and Dunn	Electrofished: 1 site, juvenile SH were seen.	1995
Bridge Creek T15N, R14W, Sec. 20	SH COH	RM-0.06 RM-?	Coyle, Chimarios and Dunn Brown and Moyle	Electrofished: 2 sites, juvenile SH were found as far as 326 feet upstream. Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1995 1991
Shingle Mill Creek T15N, R14W, Sec. 17	SH	RM-0.4	Coyle, Chimarios and Dunn	Electrofished: juvenile SH were found upstream to 1,887 feet.	1995
Mc Carvey Creek T15N, R14W, Sec. 16	SH	RM-0.4	Coyle, Chimarios and Dunn	Electrofished: 2 sites, juvenile SH were found 2,106 feet above the mouth.	1995
Rose Creek T15N, R14W, Sec. 23	SH	RM-0.6	Flosi, Coyle, Chimarios and Dunn	Electrofished: 2 sites, found juvenile SH upstream 3,330 feet.	1995
Low Gap Creek T15N, R14W, Sec. 23	SH	RM-0.85	Flosi	Electrofished: found juvenile SH 4,508 feet upstream.	1995
Soda Creek T15N, R14W, Sec. 14	SH	RM-0.04	Flosi, Chimarios and Dunn	Electrofished: 1 site, found juvenile SH 213 feet above the mouth.	1995

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
North Branch North Fork Navarro River T15N, R15W, Sec. 7	COH SH	RM-4.0 RM-4.0	Brown and Moyle Jones, Bysshe and Strayer	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. Electrofished: 1 site, found juvenile COH and SH 1,000 feet above the East Branch.	1991 1994
Cook Creek T15N, R15W, Sec. 6	SH	RM-1.7	Jones, Bysshe and Strayer	Electrofished: found juvenile SH 9,102 feet upstream.	1994
Dutch Henry Creek T16N, R15W, Sec. 32	SH	RM-?	Jones	Personal observation: have found juvenile SH in this stream.	1961
John Smith Creek T16N, R15W, Sec. 33	COH SH	RM-0.6 RM-0.6	Brown and Moyle Jones, Bysshe & Strayer	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. Electrofished: 3 sites, juvenile COH and SH were present 3,208 feet above the mouth.	1991 1994
Little North Fork North Fork Navarro River T16N, 15W, Sec. 33	SH	RM-2.9	Brown and Moyle Jones, Bysshe & Strayer	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. Electrofished: 1 site, juvenile SH found 15,559 feet upstream of the confluence with the North Branch of the North Fork Navarro River.	1991 1994
Redwood Creek T16N, R15W, Sec. 35	SH	RM-?	Jones	Personal observation: juvenile SH have been seen near the mouth of this stream. The extent of their use was unknown.	1961
Bottom Creek T16N, R15W, Sec. 36	SH	RM-0.3	Flosi, Jones, Bysshe & Strayer	Electrofished: 2 sites, juvenile SH found 1,674 feet upstream.	1994

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Mill Creek T14N, R15W, Sec. 3	SH COH	RM-? RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. A stream survey is needed.	1991
Meyer Gulch T15N, R15W, Sec. 27	?	RM-?		A stream survey is needed.	
Hungry Hollow Creek T15N, R15W, Sec. 36	SH	RM-1.0	Retherford	Personal observation: juvenile SH were present 1.0 miles upstream.	1998
Indian Creek T14N, R14W, Sec. 19	SH COH	RM-6.5 RM-?	Brown and Moyle Coyle Bowers and Nebenzahl	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. Personal observation: juvenile SH were present 6.5 miles. Snorkel survey: found juvenile SH and 2 COH present near the North Fork of Indian Creek in July. Five stations were sampled: # 1 found 100 SH YOY, 50 SH Y+, # 2 found 55 SH YOY, 20 SH Y ¹⁺ , and 16 SH Y ²⁺ , # 3 found 100 SH YOY, 7 SH Y ¹⁺ , 1 Y ²⁺ and 1 YOY COH, # 4 found 50 SH YOY, 20 SH Y ¹⁺ and 1 YOY COH, # 5 found 2 SH YOY.	1991 1994 1997
West Branch Indian Creek T14N, R14W, Sec. 8	?	RM-?		A stream survey is needed.	
Bullock Creek T14N, R14W, Sec. 5	?	RM-?		A stream survey is needed.	
Gschwend Creek T15N, R14W, Sec. 32	?	RM-?		A stream survey is needed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
North Fork Indian Creek T14N, R14W, Sec. 14	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991 A stream survey is needed.	1991
West Branch North Fork Indian Creek T15N, R14W, Sec. 35	?	RM-?		A stream survey is needed.	
Unnamed Tributary to North Fork Indian Creek T15N, R14W, Sec. 36	?	RM-?		A stream survey is needed.	
Gut Creek T14N, R13W, Sec. 18	SH COH	RM-0.1 RM-?	Brown and Moyle Marke and Fish	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. Snorkel survey: found juvenile SH present in 10 pools above the mouth of Gut Creek. The first pool had 15 SH YOY, the second pool had 14 SH YOY. Overall the number of juvenile SH varied per pool from 14 to 40, with only 1 Y+ age sized SH.	1991 1997
Anderson Creek T14N, R14W, Sec. 19	SH	RM-12.5	Michaels & Thompson Albert Coyle	Survey: two sets of falls, one 6 feet high and the other 10 feet high mark the upstream limit of SH value. Electrofished: downstream of the Rossi Bridge the juvenile SH population estimate was 133 fish/100 feet of stream. Personal observation: juvenile SH were found at 1.7 miles upstream.	1969 1970 1994

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Con Creek T14N, R14W, Sec. 34	SH	RM-0.9	Coyle Jones	Field note: Juvenile SH used this stream. Personal observation: a barrier prevents salmonids from migrating out of the valley floor into 5 miles of spawning and rearing area located in the headwater. The landowner has refused to allow removal. The habitat above the barrier is good and a resident rainbow trout population is all that remains of the anadromous use in the upper drainage.	19994 1996
Ornbaun Creek T14N, R14W, Sec. 34	SH	RM-?	Coyle	Field note: juvenile SH used this stream.	1994
Robinson Creek T13N, R14W, Sec. 2	SH	RM-2.7	Jones Albert Coyle	Personal observation: juvenile SH were present. Electrofished: found juvenile SH 6/100 feet of stream. Field note: Juvenile SH were observed in this stream.	1967 1970 1994
Mill Creek T13N, R14W, Sec. 2	SH	RM-1.5	Michaels & Thompson Anderson	Survey: two permanent dams 30 feet high, juvenile SH were present below. Electrofished: found 30 juvenile SH/ 100 feet of stream.	1969 1970
Soda Creek T13N, R14W, Sec. 2	SH	RM-3.8	Jones Coyle	Personal observation: juvenile SH were present up to a 20 foot high falls. Field note: SH use this stream.	1970 1994
Jimmy Creek T13N, R13W, Sec. 17	SH	RM-2.3	Michaels & Thompson	Survey: juvenile SH were present. The extent of their use was unknown.	1969
Rancheria Creek T14N, R14W, Sec. 19	SH COH	RM-30.5 RM-?	Moore & Keller Brown and Moyle MacMillan, Melton, D. Jones & Huber	Survey: juvenile SH were present. Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. Electrofished: 4 sites, juvenile SH were present at 11,180 feet above the mouth.	1962 1991 1994

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Ham Canyon Creek T14N, R14W, Sec. 30	COH	RM-1.4	Peters & Ayers	Survey: both juvenile COH and SH were present.	1969
	SH	RM-2.1	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			MacMillan & Onradnik	Electrofished: 2 sites, juvenile SH were present 11,180 feet upstream.	1996
Dago Creek T14N, R15W, Sec. 26	COH	RM-2.5	Peters & Ayers	Survey: juvenile COH and SH were present.	1969
	SH	RM-?	MacMillan & Huber	Electrofished: 1 site, juvenile SH were present 50 feet above the mouth.	1996
Horse Creek T13N, R15W, Sec. 1	COH	RM- 2.5	Peters & Ayers	Survey: juvenile SH were present.	1969
	SH	RM-2.5	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			MacMillan & Ouradnik	Electrofished: 1 site, juvenile SH were found 2,610 feet upstream.	1996
Bear Wallow Creek T13N, R14W, Sec. 8	SH	RM-2.0	Keller	Survey: juvenile SH were present.	1962
			MacMillan & Ouradnik	Electrofished: 2 sites, juvenile SH were found 1,786 feet upstream.	1996
Minnie Creek T13N, R14W, Sec. 17	COH	RM-1.7	Peters & Ayers	Survey: juvenile COH and SH were present.	1969
	SH	RM-1.7+	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Coyle	Field note: SH use this stream.	1994
Alder Creek T13N, 14W, Sec. 17	SH	RM-1.6	Jones	Personal observation: juvenile SH were present.	1969

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Camp Creek T13N, R14W, Sec. 17	SH	RM-6.0	Henry	Survey: juvenile SH were present.	1977
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Coyle	Field note: SH use this stream.	1994
German Creek T13N, R14W, Sec. 33	SH	RM-1.4	Jones	Personal observation: juvenile SH were present.	1962
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1991
Beasley Creek T13N, R13W, Sec. 32	SH	RM- 0.5	Keller & Moore	Survey: juvenile SH were found in this stream.	1962
			Coyle	Field note: SH use this stream.	1994
Yale Creek T12N, R13W, Sec. 14	SH	RM-0.8	Jones	Personal observation: juvenile SH have been seen in this stream.	1962
Unnamed tributary to Rancheria Creek T12N, R12W, Sec. 30	SH	RM-?	Jones	Personal observation: juvenile SH have been seen in this stream but their range was undetermined.	1978
Greenwood Creek T15N, R17W, Sec. 35	COH	RM-4.6	Netherby & Rockwood	Survey: juvenile COH and SH were present in the lower drainage up to a 20 foot barrier falls. Resident trout were present in the headwater but their range was unknown.	1966
	SH	RM-4.6			
	RT	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Elk Creek T14N, R17W, Sec. 2	COH	RM-6.9	Finlayson & Nelson	Survey: juvenile SH but no COH were observed in 1973.	1973
	SH	RM-6.9	Baracco, Charters, Moore	Survey: including electrofishing, COH and SH were present.	1976
	RT	RM-?	Jones	Personal observation: a barrier falls exists on this stream, above which a resident trout population occurs. The extent of the resident trout population is unknown.	1981
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1991
Mallo Pass Creek T14N, R17W, Sec. 36	SH	RM-1.4	Van Zandt, Jones,	Survey: barrier log jams stop migrations of anadromous salmonids. About 3 miles of stream was used at the time of the survey.	1975
	RT	RM-1.9	Jones	Electrofished: 3 sites, juvenile SH were present.	1988
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Daugherty	Electrofished: found RT in the SE quarter of section 33.	1996
Alder Creek T13N, R17W, Sec. 12	SH	RM-4.0	Netherby & Rockwood	Survey: a barrier, 20 foot high falls was present 4.0 miles upstream. Juvenile SH were present below the barrier.	1966
	RT	RM-6.0	Henry	Survey: juvenile SH were present.	1977
			Daugherty	Electrofishing: found SH/RT present. Presumed to be juvenile SH but in reality may have been RT. The sampling station was 11 miles upstream in Section 22. Another station upstream revealed no fish.	1996
Nye Creek T13N, R16W, Sec. 10	RT	RM-0.09	Daugherty	Electrofished: found juvenile RT one + age sized fish.	1995 and 1996
Tin Can Creek T13N, R16W, Sec. 11	RT	RM-0.06	Daugherty	Electrofished: found RT above the mouth 300 feet. Also found one tailed frog.	1996

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
John Creek T13N, R16W, Sec. 12	RT	RM-0.5	Daugherty	Electrofished: found RT present ½ mile upstream of the mouth.	1996
Bee Tree Creek T13N, R16W, Sec. 11	RT	RM-0.25	Daugherty	Electrofished: found RT present 1320 feet above the mouth.	1996
Unnamed Tributary to Alder Creek T13N, R16W, Sec. 12	RT	RM-0.2	Daugherty	Electrofished: found RT present 0.2 miles above the mouth.	1996
Brush Creek T13N, R17W, Sec. 23	COH	RM-9.0	CDFG	Survey: found juvenile COH and SH present in 9 miles of Brush Creek.	1968
	SH	RM-11.0	Jones	Survey: The survey included the use of electrofishing, 2 sites, 228 juvenile SH were present/ 100 feet in the upper station and 279 fish/ 100 feet in the lower station.	1981
			Kidd and Painter	Electrofished: found only juvenile SH present.	1983
			Snyder	Electrofished: the total calculated juvenile SH population was 23,978 fish per mile of stream.	1984
			Jones and Walton	Electrofished: found only juvenile SH up to and above the confluence of the South Fork.	1986
		Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1991	
Spanish Creek T13N, R16W, Sec. 19	SH	RM-1.25	CDFG	Survey: found juvenile COH present in 1.25 miles of Spanish Creek.	1968
			Jones	Survey: including electrofishing, found only juvenile SH present.	1983
Mill Creek T13N, R16W, Sec. 29	SH	RM-0.6	Nicol, Litwin and Swirhun	Survey: found were juvenile "SH, possibly COH", ranging in size from 2 to 6 inches in numbers of 25 - 50 fish per pool.	1980
	COH ?	RM-?	Jones	Survey: juvenile SH were present.	1983

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Garcia River T13N, R17W, Sec. 34	COH	RM-?	Kimsey	Memo: COH & SH were both present.	1952
	SH	RM-?	Jones, Susan, Daugherty, Wooster and Moore	Electrofished: observed 6 adult salmonids. Caught and identified 2 adult COH and 2 SH in the Garcia River, 2-1/2 miles above the Eureka Hill Road, (1/21/87).	1987
	CHIN	RM- ?	Daugherty	Electrofished: 1987- main stem above signal Creek, only juvenile SH were present.	1987
			Jones	Electrofished: found only juvenile SH.	1987
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones	Electrofished: within the mainstem above Rolling Brook juvenile SH were found. In the main stem down stream of Windy Hollow Road only juvenile SH were found. In the main stem above East End Creek only juvenile SH were present.	1993
			Harris	Electrofished: found only juvenile SH.	1996
			Daugherty	Electrofished: found juvenile SH present in Section 11 above Inman Creek.	1996
North Fork Garcia River T12N, R16W, Sec. 10	SH	RM-5.8	Johnson	Survey: juvenile SH were present, (in the upper 3 miles, 50 fish/100 feet), no COH were found.	1967
			Swirhun and Litwin	Survey: juvenile "SH were common to abundant". They ranged from 6 to 8 inches.	1980
			Jones	Electrofished: 1983, 1987, and 1990 found only juvenile SH each year.	1983, 1987, 1990
Unnamed Tributary to Garcia River T12N, R16W, Sec. 25	SH	RM-0.09	Daugherty	Electrofished: found juvenile SH present 500 feet upstream of the mouth.	1996

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Rolling Brook Creek T12N, R16W, Sec. 24	SH	RM-1.0	Jones Daugherty	Electrofished: juvenile SH were present. Electrofished: found juvenile SH present about ½ mile above the mouth.	1987 1996
South Fork Garcia River T12N, R15W, Sec. 30	COH SH	RM-2.0 RM-2.0	Jones Daugherty	Electrofished: 1986, juvenile SH were present, 1987 juvenile COH and SH, 1988 juvenile COH and SH, 1989, 1990, 1991 and 1992 only juvenile SH. Electrofished: found juvenile COH and SH present in 1994, in 1995 only juvenile SH, in 1996 juvenile COH and SH were both present. The site was 500 feet above the mouth.	1986, 1987, 1989, 1990, 1991, 1992 1994, 1995, 1996.
Fleming Creek T11N, R15W, Sec. 4	COH SH	RM-0.1 RM-?	Jones Daugherty	Electrofished: 1987 juvenile SH were present, 1988 COH & SH, 1989 1990 and 1991 only SH were present. Electrofished: found juvenile SH present 0.4 miles upstream in July.	1988, 1989, 1990, 1991 1996
Signal Creek T12N, R15W, Sec 26	SH	RM-0.6	Jones Maahs	Electrofished: juvenile SH were present. Carcass survey: one redd was found, 3 adult SH were also found.	1987, 1995 1996
Inman Creek T12N, R15W, Sec 14	COH SH	RM-3.8 RM-3.8	Daugherty Jones Maahs	Electrofished: found juvenile COH and SH present at the mouth in 1990. Also found juvenile SH present in 1996 Electrofished: found only juvenile SH present. Carcass survey: only one COH carcass was found.	1990 1996 1996 1997
Blue Water Hole Creek T12N, R15W, Sec. 11	SH	RM-1.8	Jones	Personal observation: juvenile SH were present above the mouth of the stream.	1988
Larmour Creek T12N, R14W, Sec. 5	SH	RM-1.3	Jones	Survey: juvenile SH were present but a 75 foot falls, 1.3 miles above mouth blocks further upstream SH migration.	1975

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Mill Creek T12N, R13W, Sec. 7	SH	RM-5.5	Holman Maahs Maillaird	Survey: juvenile SH were present. Carcass survey, 2 adult SH were seen. Personal observations: the land owner revealed the presence of SH 5.5 miles upstream.	1968 1996 1999
Redwood Creek T12N, R13W, Sec. 6	SH	RM-1.5	Holman	Survey: juvenile SH were present but the stream was considered of poor value.	1968
Pardaloe Creek T12N, R13W, Sec. 7	SH	RM-2.3	Jones Jones Maahs	Personal observation: saw juvenile SH in this stream. Electrofished: found only juvenile SH. Carcass survey: two adult SH were seen.	1975 1994 1996
Point Arena Creek T12N, R17W, Sec. 14	SH	RM-1.7	Jones	Personal observations: including electrofishing, found juvenile SH in the area of stream within the town of Point Arena.	1975
Moat Creek T12N, R16W, Sec. 30	none	RM-?	Jones	Electrofished: found no fish.	1986
Ross Creek T12N, R16W, Sec. 29	?	RM-?		A stream survey is needed.	
Galloway Creek T12N, R16W, Sec. 29	none	RM-?	Nicol and Litwin	Survey: no fish were observed. Found was a "massive" log jam considered to be a "complete barrier" at the mouth of the stream. This jam was reported to be 150 feet long, 100 feet wide and 10 feet high.	1980

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Schooner Gulch T12N, R16W, Sec. 32	SH	RM-2.0	Michaels & Thompson	Survey: juvenile SH were present.	1969
	COH	RM-?	Jones and Walton	Electrofished: found juvenile SH present.	1986
			Cox	Electrofished: juvenile SH were present.	1990
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Daugherty	Electrofished: found juvenile SH in Section 34, 2miles above the mouth.	1995 and 1996
North Fork of Schooner Gulch T12N, R16W, Sec. 29	SH	RM-1.4	Michaels & Thompson Brown and Moyle	Survey: juvenile SH were present. Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1969 1991
Hall Creek T12N, R16W, Sec. 34	SH	RM-0.04	Daugherty	Electrofished: found juvenile SH present 200 feet upstream.	1995 and 1996
Fish Rock Gulch T11N, R15W, Sec. 18	SH	RM-?	Jones	Personal observation: the culvert under Hwy 1 had a fish passage problem in 1997 and needed work. This work was never completed.	1997-2000
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Harris	Electrofished: found juvenile SH present 200 yards above the problem culvert.	1999
Gualala River T11N, R15W, Sec. 27	SH	RM- 3.4	Jones	Personal observation: juvenile and adult SH were present.	1986,1995, 1996
	COH	RM-3.4	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
North Fork Gualala River T11N, R15W, Sec. 23	COH	RM-12.5	Pool & Parker	Survey: both COH and SH were present.	1964
	SH	RM-12.5+	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones	Electrofished: only juvenile SH were present. CDFG planted yearling COH in the Little North Fork in 1995, 1996 and 1997.	1986,1995, 1996, 1997
Little North Fork Gualala River T11N, R15W, Sec. 23	COH	RM-4.0	Parker	Survey: both juvenile COH and SH were present.	1964
	SH	RM-4.0+	Jones	Electrofished: 1983 and 1986, juvenile SH were present. In 1988 juvenile COH and SH were both present. In 1989 through 1993 and again in 1995, only juvenile SH were found.	1983, 86, 88, 89, 90, 91, 92, 93, 95
			CDFG	Planted COH three years (1995, 96 and 97).	1995-96-97
Doty Creek T11N, R15W, Sec. 10	SH	RM-1.0	Jones	Electrofished: juvenile SH were present.	1986
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones and Harris	Electrofished: spot checking, revealed the presence of only juvenile SH.	1995
Robinson Creek T11N, R15W, Sec 12	SH	RM-1.0	Cox	Survey: juvenile SH were seen.	1980
			Jones	Electrofished: juvenile SH were present.	1983, 1986
Unnamed Tributary to North Fork Gualala River T11N, R14W, Sec. 7	SH	RM-1.8	Jones	Personal observation: juvenile SH were present but fishery value ended at a six foot falls.	1986
Dry Creek T11N, R14W, Sec. 6	SH	RM-1.2	Jones	Electrofished: juvenile SH were present.	1986, 1994

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Stewart Creek T11N, R14W, Sec. 9	SH	RM-0.8	Jones	Personal observation: juvenile SH were present and their use ended at a 15 foot falls 0.8 miles upstream.	1986
Billings Creek T12N, R14W, Sec. 34	SH	RM-6.5	Jones	Personal observation: juvenile SH were seen in this stream.	1986
Robinson Creek 12N, R14W, Sec. 34	SH	RM-2.0	Jones	Personal observation: juvenile SH were present in this stream.	1986
South Fork Gualala River T11N, R15W, Sec. 26				See Sonoma County	
Rockpile Creek T11N, R14W, Sec. 31	SH	RM-?	Jones Parker	Personal observation: The mouth of this stream and about 6 miles are within Sonoma County. An additional 14 miles are within Mendocino County. Described below is the fishery value of the Mendocino County portion. Survey: found juvenile SH present in densities of 100/100 feet of stream, 1 inch to 1 foot long. The survey extended through both counties for a distance of 20 miles. The range of juvenile SH observation in Mendocino County was not listed in the survey.	2000 1964
Unnamed Tributary to Rockpile Creek T11N, R13W, Sec. 6	SH	RM-0.06	Parker	Survey: found a small population of juvenile SH below the barrier (located 100 yards above the mouth). This stream. . ." very little fisheries value because of low flow and dry conditions during the summer."	1964
Unnamed Tributary to Rockpile Creek T11N, R13W, Sec. 4	SH	RM- 2.0	Greene	Survey: found juvenile RT/SH present , 1-15 inches and averaging 3 inches present. Species were not specified, but will assume they were juvenile SH.	1964

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Big Pepperwood Creek T11N, R14W, Sec. 31	SH	RM-4.0	Rowell Jones	Survey: juvenile SH were present 3 to 8 inches in length 6 per pool in the upper reaches in September. Personal information: tributary to South Fork Gualala River with confluence in Sonoma County. The upper 2.4 miles of fishery value is in Mendocino County.	1964 1997
Dry Creek Mendocino County, tributary to Lake Sonoma in Sonoma County, hence Russian River T11N, R12W, Sec. 11 at Co. line.	RT	RM-12.5	Jones	Personal information: Resident RT use this stream.	1997
McChristian Creek T11N, R12W, Sec. 11	?	RM-?		A stream survey is needed.	
Franks Canyon T11N, R12W, Sec. 11.	?	RM-?		A stream survey is needed.	
Soda Creek T11N, R12W, Sec. 2	?	RM-?		A stream survey is needed.	
Big Foot Creek T12N, R12W, Sec. 27	?	RM-?		A stream survey is needed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Ingram Creek T12N, R12W, Sec. 14	?	RM-?		A stream survey is needed.	
Elkhorn Creek T12N, R12W, Sec. 16	?	RM-?		A stream survey is needed.	
MattoleRiver (the downstream portion is in Humboldt County. The following portion is in Mendocino County T5S, R2E, S34)	COH SH	RM-3.5 RM-3.5+	Jones	Electrofished: both juvenile COH and SH were present in the upper portion of the drainage.	1986
Thompson Creek T5S, R2E, Sec. 27	COH SH CHIN	RM-1.4 RM-1.4 RM-1.4	Jones	Personal observations: including electrofishing, two stations electrofished, juvenile SH were present at 0.7 miles (in Mendocino County). An additional 0.7 miles of stream in Humboldt County provides habitat for juvenile SH. CHIN, COH, and SH all use this stream.	1986
South Branch Thompson Creek T5S, R2E, Sec. ?	COH SH	RM-? RM-?	Jones	Electrofished: juvenile COH and SH were present.	1986
McNasty Creek T5S, R2E, Sec. 33	COH SH	RM-? RM-?	Jones	Electrofished: juvenile COH and SH were present.	1986

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Eel River (the following portion is in Mendocino Co.)	COH	RM-30.5 (in Mendo. Co.)	Jones	Personal observation: COH migrate in the main stem Eel River up to Outlet Creek. During the 1946-47 trapping period 47 COH salmon were trapped at Van Arsdale.	1983
	SH	RM-71.0 (in Mendo. Co.)	Jones & Thompson	Survey: completed from Scott Dam to Van Arsdale, juvenile SH were observed throughout.	1969
	SSH	RM-71.0 (in Mendo. Co.)	Doo, Michaels & Anderson	Survey: from Van Arsdale to 2.5 mi. below Hurst, juvenile SH were observed.	1969
	CHIN	RM-71.0 (in Mendo. Co.)	Meyers, Thompson & Jones Puckett & Hinton	Survey: Hearst to Outlet Creek, found adult & juvenile SH present.	1969
				Survey: Hearst to Outlet Creek, found adult & juvenile SH present.	1973
				Snorkel survey: Scotts Dam to Van Arsdale, counted 1 SSH but 19 others were reported by Mulligan the CDF&G person at Van Arsdale, as present in front of the PGE screen. Later in the summer the presence of these fish was confirmed.	1985
	RT	RM-2.2	Walton Jones & Baker	Creel census: Dos Rios to Outlet Creek, 12/2/86, 78 anglers caught 37 CHIN during a total of 353.75 hours. 12/3/86, 49 CHIN were caught in 356.75 hours. On 12/13/86, 71 anglers caught 10 CHIN in 195.75 hours.	1986
				Survey: canoed from Dos Rios to Alder Point, saw juvenile SH the entire distance.	1986
				Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
	Murphy & Rodriguez	Grass	Survey: the origin of the Eel River is in Mendocino County and flows into Lake County above Lake Pillsbury and then back into Mendocino County, hence on to Trinity County, Humboldt County and on to the Ocean. The upper 3.2 miles of the headwaters flows through extensive barriers, however resident rainbow trout were found throughout this upper section.	1993	
CDFG records: One COH was trapped at the Van Arsdale Fishery Station during the 2000/2001 season. salmonid trapping data is available from CDF&G. Trapping records extend from 1933/34.			2001		

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
South Fork Eel River (in Mendocino County)	COH SH SSH? CHIN	RM-54.5 RM-54.5 RM-? RM-54.5	Jones	Personal observation: snorkel and carcass surveys have found COH, SH and CHIN in this stream. Summer steelhead were once present in the South Fork.	1986
East Branch South Fork Eel River T4S, R3E, Sec.36	SH SSH? CHIN	RM- 19.9 RM-? RM?	Jones & Johnson Coyle, Mesman, MacDownell, Goodfield,& Mitchell.	Personal observation: have seen large salmonids in pools in the headwater during the summer, suspected summer steelhead, but close observations were impossible. CHIN are also known to use this stream but their range has not been determined. A stream survey is needed. Electrofished: 5 sites, juvenile SH were present 105,085 feet above the mouth.	1982 1993
Elkhorn Creek T24N, R16W, Sec. 10	SH CHIN	RM-2.0? RM?	Cherr & Griffin	Stream inventory: indicates RT were present. A stream survey is needed.	1979
Foster Creek T 24N, R16W, Sec. 16	SH	RM-0.6	Cherr & Griffin	Stream inventory: indicates RT were present. A stream survey is needed.	1979
Cruso Cabin Creek T24N, R16W, Sec. 10	SH?	RM-?		Unknown fishery resource. A stream survey is needed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Low Gap Creek T5S, R3E, Sec. 24	SH	RM-1.8	Wooley	Survey: A 40 foot falls blocks further migration, but juvenile SH were present below the falls.	1980
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Coyle & Mitchell	Electrofished: 2 sites, juvenile SH were found 5,988 feet above the mouth.	1993
Indian Creek T5S, R3E, Sec. 35	SH	RM-9.5	Ayers & Peters	Survey: a 15 foot falls limits anadromous migrations. Juvenile SH were present below.	1968
	COH	RM-8.2	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Michaels, Mesman & Coyle	Electrofished: 2 sites, juvenile SH and COH were present to the upper station a distance of 43,306 feet, (8.2 miles).	1993
Jones Creek T5S, R3E, Sec. 27	COH	RM-0.1	Jones	Electrofished: juvenile COH and SH were confirmed present.	1993
	SH	RM-0.8	Coyle & Michaels	Electrofished: 2 sites, both juvenile COH and SH were present in the lower site 519 feet above mouth. Juvenile SH were also present in the upper site.	1993
Parker Creek T5S, R3E, Sec. 28	SH	RM-0.2	Jones	Personal observation: juvenile SH were present but a 25 foot falls blocked further upstream migration.	1968
Moody Creek T24N, R18W, Sec. 5	SH	RM-1.4	Jones	Electrofished: juvenile SH were present.	1987
	COH	RM-1.4	Mesman & Michaels	Electrofished: 3 sites, juvenile COH and SH were present 1,727 feet upstream.	1993
Sebbas Creek T24N, R18W, Sec. 5	SH	RM-2.3	Jones	Personal observation: juvenile SH were present.	1968
			Michaels & Mesman	Electrofished: 3 sites, juvenile SH were present 12,004 feet above the mouth.	1993

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Coulborn Creek T24,N, R18W, Sec. 6	SH	RM-1.2	Jones	Personal observation: juvenile SH were present.	1968
	COH	RM-1.1	Michaels, Mesman & Elliot	Electrofished: 3 sites, Juvenile COH and SH were found 5,690 feet above the mouth but only juvenile SH were present 6,201 feet above the mouth.	1993
Anderson Creek T24N, R18W, Sec. 7	SH	RM-4.6	Jones	Personal observation: juvenile SH were present.	1968
	COH	RM-4.6	Coyle & Mesman	Electrofished: 1 site, both juvenile COH and SH were present 24,296 feet above the mouth.	1993
Piercy Creek T5S, R3E, Sec. 35	SH	RM-4.2	Peters & Ayers	Survey: juvenile COH and SH were present.	1968
	COH	RM-2.0	Boccone & Rowser	Survey: found a logging road barrier in the headwater with a 12 foot falls. Juvenile SH were present below.	1977
			Fellman, Litwin and Swirhun	Survey: juvenile SH were abundant, 150/ 100 feet of stream.	1980
			Sortori & Long	Electrofished: 5 sites, juvenile COH and SH were present 10,317 feet, (2.0 miles) upstream, but only juvenile SH were present 13,611 feet (2.6 miles) above the mouth.	1990
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1991
Stanley Creek T24N, R18W, Sec. 1	SH	RM-4.0	Jones	Personal observations: juvenile SH were present.	1968
	COH	RM-0.1	Curtis & Jones	Carcass survey: found CHIN carcasses.	1988
	CHIN	RM-?	Madigan and Mizuno	Stream survey: found one live CHIN and one carcass.	1990
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Elliot, Monroe and Checkler	Electrofished: 4 sites, juvenile COH and SH were present 775 feet upstream.	1992

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
McCoy Creek T24N, R17W, Sec. 6	COH	RM-3.5	Peters & Ayres	Survey: juvenile COH and SH were present, nursery habitat was poor.	1968
	SH	RM-3.5	Mullins, Humpreys & Goodfield	Electrofished: 1 site, found juvenile SH present 244 feet upstream.	1995
North Fork McCoy Creek T24N, R17W, Sec. 4	COH	RM-2.4	Peters & Ayres	Survey: juvenile COH and SH were present, nursery habitat was considered poor.	1968
	SH	RM-2.4	Downey & Goodfield	Survey: Observed juvenile SH through the survey reach of 4,106 feet.	1995
Bear Pen Creek T24N, R17W, Sec. 7	SH	RM-4.0	Jones	Personal observation: juvenile SH were present.	1968
	COH	RM- 0.03	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Elliot & Monroe Harris	Electrofished: 1 site, found juvenile SH and COH 158 feet above the mouth. Personal observation: found only juvenile SH.	1992 1999
Cub Creek T24N, R17W, Sec. 18	COH	RM-0.2	CDFG	Resource map: COH and SH use 0.2 miles of stream.	1979
	SH	RM-0.2	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1991
Red Mountain Creek T24N, R17W, Sec. 17	SH	RM-4.2	Thompson & Jones	Survey: juvenile SH were present. Local residents indicate that when the Cedar Creek hatchery was in operation COH, (pre 1964) used the stream, (W. Jones).	1968
	COH?	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Goodfield & Robertson	Electrofished: 2 sites, juvenile SH were found 3,825 feet above the mouth.	1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Wildcat Creek T24N, R17W, Sec 20	COH	RM-3.0	Ayers & Peters	Survey: COH and SH were present in “good supply” in the lower 3/4 miles.	1968
	SH	RM-3.0	Dixie	Carcass survey: found 1 CHIN carcass, 5 skeletons and 3 live salmon that were unidentified.	1983
			Jones	Electrofished: juvenile COH and SH were present in the stream.	1987
			Brown and Moyle	Report: COH were listed in this stream in the report; “Status of Coho Salmon in California”, 1991.	1991
			Humphrey & Elliot	Electrofished: 1 site, juvenile SH were present 572 feet upstream.	1992
Bridges Creek T24N, R17W, Sec. 21	SH	RM-3.1	Fender, Swarhun & Litwin	Survey: juvenile SH were seen in “good abundance” in the upper area (In densities of 200 fish /100 ft of stream).	1980
			Goodfield & Abel	Electrofished: 3 sites, found juvenile SH present 16,572 feet upstream.	1994
			Coyle, Chimarios &Dunn	Survey: included electrofishing as part of a habitat typing project, juvenile SH were present.	1995
Mill Creek T24N, R17W, Sec. 23	SH	RM-0.3	Ayers & Peters	Survey: found juvenile SH. However, a barrier with a falls, stopped SH migrations 0.3 miles upstream.	1968
	COH	RM-0.02	Monroe & Cleckers	Electrofished: 2 sites, found juvenile COH and SH 123 feet upstream and juvenile SH again at 1,211 feet upstream.	1992
			Jones	Personal observation: the CCC removed the barrier but the bedrock falls remains.	1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Hollow Tree Creek T23N, R17W, Sec. 10	COH	RM-18.0	Ayres & Peters	Survey: juvenile COH and SH were present.	1968
	SH	RM-18.0	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
	CHIN	RM-18.0	Jones	Electrofished, carcass surveys and personal information: electrofished for juveniles in 1983 and 1986. From 1983 through 1995, juvenile COH and SH were both present in the upper drainage. Carcass surveys in the winter revealed: CHIN, COH and SH were present. The Salmon Restoration Association maintains a CHIN trapping and spawning facility on the stream. Records are available.	1983-2000
			Harris	Electrofished: found juvenile COH and SH present each year.	1998-99
South Fork Hollow Tree Creek T23N, R17W, Sec. 29	SH	RM-0.25	Swirhun and Montoya	Survey: juvenile SH numbering 70/100 feet of stream below the 25 foot high barrier falls. The barrier was located 0.25 miles upstream of the mouth. Above the barrier was an RT population of 2 - 3 fish/ 100 feet of stream.	1981
	RT	RM-?	Jones	Personal observation: two barriers near the mouth, one about 200 feet above the mouth, the second, 20 feet high some 0.2 miles upstream. Resident fish were present above the barrier, their range was undetermined.	1997
Mule Creek T23N, R17W, Sec. 17.	RT	RM-0.5	Swirhan and Montoya	Survey: only "one juvenile SH/RT" was found during the survey. Slow velocity and stagnet pools. The survey indicates that 3 barriers were found but lists the height as 2 and 3 feet.	1981
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones	Personal observation: the fish seen by Swirhan and Montoya and reported by Brown and Moyle had to be a RT or something else as there is a 25 foot high vertical falls on the South Fork of Hollow Tree Creek below the confluence with Mule Creek (W. Jones).	2000

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Middle Creek T23N, R17W, Sec. 29	COH SH	RM-0.09 RM-0.09	Montoya and Swirhun Monroe & Cleckler	Survey: no fish observed. However, did find a falls of about 40 feet in height located about 100 yards upstream of the mouth. Electrofished: 3 sites, juvenile COH and SH were present 470 feet above the mouth. A bedrock falls 38 feet in height was found 715 feet above the mouth.	1981 1992
Islam John Creek T23N, R17W, Sec. 33		RM-0	Montoya and Swirhun Jones	Survey: found two falls of 5 and 8 feet respectively at the mouth of the stream. No fish were found. Personal observation: an 8 foot high falls near the mouth was considered a total barrier.	1981 1983
Lost Man Creek T23N, R17W, Sec. 33	SH	RM-33 feet ?	Swirhun Monroe & Cleckler	Survey: the mouth of the stream has a series of three bedrock falls of 6, 10 and 8 feet high respectively. Each was considered to be a barrier. Survey: juvenile SH were seen 33 feet upstream.	1981 1992
Walters Creek T22N, R17W, Sec. 3	SH COH	RM-? RM-?	Montoya and Swirhun Brown and Moyle	Survey: found juvenile SH present in this stream. Also found 8 barriers of wood debris from 4 to 15 feet high. The highest barrier, 15 feet high was within 0.1 miles of the mouth. Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991	1981 1991
Lost Pipe Creek T22N, R17W, Sec. 3	none	RM-0	Montoya and Swirhun	Survey: no fish were found. Four barriers of 5 to 7 feet in height were found. The first was 0.1 miles above the mouth.	1981
Bear Creek T22N, R17W, Sec. 4	None	RM-0	Montoya and Swirhun	Survey: no fish were seen. Found was a log jam barrier 12 feet high just above the mouth. Immediately above this barrier was a second one of bedrock 12 feet high.	1981

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Redwood Creek T22N, R17W, Sec. 9	COH	RM-1.0	Ayers & Peters	Survey: juvenile COH and SH were “found in abundance”.	1968
	SH	RM-1.0	Jones	Electrofished: juvenile COH and SH were present.	1983
	CHIN	RM-?	Jones and Walton	Carcass survey: found CHIN and COH present in this stream both years.	1987-88 and 1988-89
			Brown and Moyle	Report: COH were listed in this stream in the report; “Status of Coho Salmon in California”, 1991.	1991
			Elliot & Humprey	Electrofished: 4 sites, found juvenile COH and SH present 3,000 feet upstream.	1991
South Fork of Redwood Creek T22N, R17W, Sec. 8	COH	RM-1.3	Ayres & Peters	Survey: juvenile COH and SH were present.	1968
	SH	RM-?	Kidd and Swirhun	Survey: juvenile COH were present in the stream up to a log barrier 5 feet high located 0.6 miles above the mouth.	1981
Bond Creek T22N, R17W, Sec. 15	COH	RM-2.7	Ayres & Peters	Survey: only SH were present but big log jams may have stopped the COH.	1968
	SH	RM-2.7+	Ellison & Carmine	Survey: juvenile SH were present.	1980
			Jones	Electrofished: juvenile SH were present.	1983
			Jones	Electrofished: 1987 through 1990: juvenile COH and SH were present each year.	1987, 1988, 1989, 1990
			Brown and Moyle	Report: COH were listed in this stream in the report; “Status of Coho Salmon in California”, 1991.	1991
			Elliot & Humpreys	Electrofished: 3 sites, found juvenile SH 6,117 feet upstream.	1991

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Michaels Creek T22N, R17W, Sec. 14	SH	RM-2.0	Ayres & Peters	Survey: juvenile SH were present.	1968
	COH	RM-?	Montoya & Swirhun	Survey: Both juvenile COH and SH were present.	1981
			CCC's	Observation: juvenile COH and SH were observed near the mouth.	1988
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Elliot & Humpreys	Electrofishing: 3 sites, juvenile SH were observed 6,070 feet upstream.	1991
			Kusera	Personal observation: found live adult COH present.	1992/93
Lynch Creek T22, R17W, Sec. 14	COH	RM-0.5	Montoya and Swirhun	Survey: juvenile COH and SH were present up to a point below the confluence with the unnamed tributary.	1981
	SH	RM-0.5			
Waldron Creek T22N, R17W, Sec. 14	COH	RM-1.7	Swirhun	Survey: no fish were observed.	1981
	SH	RM-1.7	Jones	Electrofishing: in 1988 found juvenile COH and SH,. In 1989, 90 and 91 only juvenile SH were found.	1988- 89- 90- 91
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Huckleberry T22N, R17N, Sec. 23	COH	RM-0.8	Montoya and Swirhun	Survey: found only juvenile SH.	1981
	SH	RM-1.2	Jones	Electrofished: juvenile COH and SH were present.	1983
			Halzerland and Friedrick	Electrofishing: 3 sites, found juvenile COH and SH 4,440 feet upstream. Juvenile SH were present 6,110 feet upstream.	1990
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones	Carcass survey: two live COH were found. Carcass survey: two surveys, no carcasses or fish were observed in either.	1992/93 1993/94
Butler Creek T22N, R17W, Sec. 25	COH	RM-2.0	Jones	Electrofished: juvenile COH and SH were present.	1983
	SH	RM-2.0	Halzerland and Fredrick	Electrofished: 5 sites, found juvenile COH and SH 6,440 feet, (1.2 miles) upstream.	1990
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991/92
			Jones	Carcass survey: found one live COH.	1991
Mitchell Creek T22N, R17W, Sec. 25	SH	RM-0.6	Ayres and Peters	Survey: only juvenile SH were found. A lack of flow may limit the upstream migration in this stream.	1968

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³	
Cedar Creek T23N, R17W, Sec. 14	SH	RM-9.0	Ayres and Peters	Survey: juvenile SH were present, however, salmonids found high in the drainage, above 5 miles were believed to be resident rainbow.	1968	
	CHIN	RM-2.0	Jones, Johnson and Montoya	Survey: An 800 foot culvert with perpendicular weirs throughout was present under the Hwy 101 crossing. Juvenile SH were found ranging from 1 to 10 inches in an abundance of about 300 fish/100 feet of stream. COH were said to have once been present but were not observed during this survey.	1981	
	COH	RM-?			Walton and Gilroy	Carcass survey: found 22 CHIN carcasses, no COH.
				Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
				Goodfield & Coyle	Electrofished: juvenile SH were found 43, 434 feet (8.2 mi.) upstream.	1993
				Jones	Personal observations: Prior to 1964 a CDFG COH hatchery existed on the east side of the South Fork Eel River near the mouth of Cedar Creek. This hatchery derived its source of water from Cedar Creek. COH were present in Cedar Creek at that time but are not at this time. Cedar Creek has a near excellent supply of cool summer water. I have also found chinook carcasses above the HWY 101 culvert.	1997
North Fork Cedar Creek T24N, R16W, Sec. 32	SH	RM- 1.0	Ayres & Peters	Survey: juvenile SH were present.	1968	
Big Dan Creek T23N, R17W, Sec. 14	SH	RM-3.0	Ayers & Peters	Survey: juvenile SH were present up to a falls 8 foot high located about 1/4 mile above HWY 101.	1968	
Little Dan Creek T23N, R17W, Sec. 13	none	RM-0	Jones & Grass	Personal observation: a 60 foot falls was present near the mouth, also the HWY 101 culvert was not baffled and was also believed to be a barrier.	2000	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Low Gap Creek T23N, R16W, Sec. 19	SH	RM-1.3	Sprehn	Survey: 2 unidentified carcasses were found in March. Numerous log jam barriers were also present.	1979
	COH	RM-1.0	Fredrick & Holzerland	Electrofished: 5 sites, juvenile COH were found 5,350 feet upstream. Juvenile SH were present up to 6,798 feet.	1990
			Jones	Electrofished: found juvenile COH and SH present in October.	1990
			Flosi	Electrofished: 6 young of the year age sized SH were found in an area containing two runs and a riffle.	1995
Rattlesnake Creek T23N, R16W, Sec. 20	COH	RM-?	Ayers and Peters	Survey: juvenile COH and SH were observed.	1968
	SH	RM-9.6	Jones, Flosi & Gilroy	Carcass survey: found CHIN present.	1987/88
	CHIN	RM-?	Jones	Personal observation: including carcass surveys, CHIN were observed in the stream during many years of survey.	1989
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Goodfield & Mitchell	Electrofished: 2 sites, juvenile SH were present 26,960 feet upstream.	1993
		Harris	Carcass survey: no fish were found during the 1999/00 season.	1999/00	
Foster Creek T23N, R16W, Sec. 22	SH	RM-2.5	Goodfield & Mitchell	Electrofished: 2 sites, juvenile SH were present 8,110 feet upstream.	1993
	CHIN ?	RM-?	Jones	Personal observation: SH were present. CHIN may use this stream as well.	1995
Elk Creek T23N, R16W, Sec. 23	SH	RM-1.8	Klamt, Bruno	Survey: SH were present, a fishway was located near the mouth.	1971
			Litwin, Fender & Swirhun	Survey: small population of juvenile SH were seen.	1980
			Goodfield & Mitchell	Electrofished: 2 sites, found juvenile SH present as far as 3,151 feet upstream.	1993

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Cummings Creek T23N, R15W, Sec. 19	COH	RM-0.8	Ayres, Peters	Survey: COH and SH were present, up to the 30 foot falls.	1968
	SH	RM-0.8	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Goodfield & Mitchell.	Juvenile SH were present at 942 feet upstream. A barrier was present at 2,525 feet upstream.	1993
Twin Rocks Creek T23N, R15W, Sec. 20	SH	RM-2.0	Ayers, Peters	Survey: juvenile SH were present up to the 40 foot falls.	1968
			Fender, Litwin & Swirhun	Survey: juvenile SH were present.	1980
			Goodfield & Mitchell	Electrofishing: 2 sites, found juvenile SH upstream to 913 feet.	1993
Grapewine Creek T23N, R15W, Sec. 28	SH	RM-0.75	Pinkham, Jones	Survey: including electrofishing, found 148 juvenile SH/ 100 feet. A dam built illegally on this stream cut off 1 mile of stream to anadromous fish use. Population estimates based on survey results and electrofishing revealed an estimated 5,861 juvenile SH below the dam and 4,095 juvenile resident RT above the dam.	1976
			Goodfield & Mitchell	Electrofished: 1 site, found juvenile SH 87 feet above the mouth.	1993

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Ten Mile Creek T22N, R16W, Sec. 16	COH	RM-8.0	Peters	Survey: juvenile COH and SH were present up to the Big Rock Creek tributary.	1969
	SH	RM-19.0	Swirhun, Fender and Litwin	Survey: juvenile SH present. October water temperatures ranged from 50 °F in the upper reach surveyed to 56 °F in the lower area.	
	CHIN	RM-?	Reynolds	Carcass survey: CHIN were observed.	1980
			Leos	Carcass survey: 13 surveys, CHIN were found, 52 carcasses and 12 skeletons.	1982
			Jones	Personal observation: CHIN were found up to 16 miles , on carcass surveys.	1983
			Jones	Carcass survey: CHIN and COH were present.	1986
			Jones and CCC's	Carcass survey: found 373 CHIN carcasses.	1987/88
			Goodfield & Humphrey	Electrofished: 3 sites, found juvenile SH present 52,587 feet above the mouth.	1988/89
		Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991	
Grub Creek T22N, R15W, Sec. 18	SH	RM-2.0	Jones	Personal observation: juvenile SH were present near the mouth.	1967
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
Streeter Creek T22N, R15W, Sec. 22	SH	RM-3.1	Jones	Carcass surveys: SH, COH and CHIN were present.	1987/88
	CHIN	RM-0.5+	Jones	Carcass survey: found CHIN present.	1988/89
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Goodfield & Abel	Electrofished: 4 sites, found juvenile SH present 16,627 feet above the mouth.	1994
Lewis Creek T22N, R15W, Sec. 27	SH	RM-1.0	Peters	Survey: found juvenile SH. This stream provides good nursery habitat.	1969
			Goodfield & Abel	Electrofished: 3 sites, juvenile SH were present 4,576 feet upstream.	1994

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Big Rock Creek T22N, R15W, Sec. 27	COH	RM-2.5	Peters	Survey: juvenile COH and SH were present in good numbers.	1969
	SH	RM-2.5+	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Goodfield & Abel	Electrofished: Juvenile COH were present 1,488 feet upstream. Juvenile SH were present 10,526 feet upstream.	1994
			Harris and Ingbar	Personal observation: observed a CHIN carcass in February.	2000
Wilson Creek T22N, R15W, Sec. 35	SH	RM-1.7	Peters	Survey: found juvenile SH. Low flows appears to be a limiting factor.	1969
	CHIN	RM-?	Cal Trans	Field note: CHIN were once blocked at the HWY 101 crossing but this passage problem has been corrected.	1991
			Ingram	Personal observation: a high falls was found just south of Hwy 101, no fish were observed.	1993
Mud Springs Creek T22N, R15W, Sec. 11	COH	RM-?	Cherr & Griffin	Fisheries inventory: COH and SH were present.	1979
	SH	RM?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
Little Case Creek T22N, R15W, Sec. 11	SH-?	RM-?		A stream survey is needed.	
Mill Creek T21N, R15W, Sec. 11	SH?	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
	COH	RM-?			

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Cahto Creek T21N, R15W, Sec. 12	COH	RM- 1.7	Cherr & Griffin	Stream inventory: both COH and SH were present.	1979
	SH	RM-3.0	Jones	Personal observation: CHIN and SH were present. However, COH were not found.	1986
	CHIN	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones	Electrofished: found only juvenile SH.	1993
			Goodfield, Kramer & Garrett	Electrofished: 2 sites, juvenile SH were found 15,640 feet upstream.	1993
Unnamed Tributary to the South Fork Eel River T22N, R16W, Sec. 20	SH	RM-0.2	Jones and Montoya	Survey: juvenile SH were present for 0.2 miles.	1981
Fox Creek T22N, R16W, Sec. 21	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
McKinley Creek T22N, R16W, Sec. 21	?	RM-?		A stream survey is needed.	
Elder Creek T22N, R16W, Sec. 29	SH	RM-3.3	Hunt	Survey: SH were present.	1975
	CHIN	RM-?	Reynolds	Field note: the caretaker at the Nature Conservancy preserve, reported that he saw a single CHIN in the mouth of Elder Creek.	1982
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Flosi & Downey	Personal observation: juvenile SH were found 8,601 feet upstream.	1992

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Jack of Hearts Creek T22N, R16W, Sec. 29	SH	RM-2.0	Jones	Personal observation: juvenile SH were present 2.0 miles upstream.	1968
	COH	RM-0.5+	Fender, Litwin and Swirhun	Survey: juvenile COH and SH were present in numbers of 20/100 feet of stream, up to a barrier 15 feet high. The barrier was located about 2.0 mile upstream.	1980
	CHIN	RM-0.5+	Walton and Long	Carcass survey: found CHIN and COH present.	1987/88
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Davis & Mitchell	Electrofished: 2 sites, juvenile SH were present 10,942 feet upstream.	1992
Deer Creek T21N, R16W, Sec. 4	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
Little Charlie Creek T21N, R16W, Sec. 9	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
Dutch Charlie Creek T21N, R16W, Sec. 9	CHIN	RM-3.5	Sprehn, Clark	Survey: 6 salmon were seen in February.	1979
	SH	RM-3.5 +	Leos	Carcass survey: 20 CHIN carcasses and 7 skeletons were found.	1983
	COH	RM-1.7	Jones	Personal observation: juvenile SH were present.	1986
			Jones and Curtis	Carcass survey: found both CHIN and COH present.	1987/88
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Crittendon & Mitchell	Electrofished: 3 sites, juvenile COH and SH were present 8,729 feet upstream.	1992
Thompson Creek T21N, R16W, Sec. 8	?	RM-?		A stream survey is needed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Eagle Creek T21N, R17W, Sec. 12	?	RM-?		A stream survey is needed.	
Redwood Creek T21N, R16W, Sec. 16	COH	RM-2.3	Leos	Carcass survey: 4 CHIN were seen in January.	1983
	CHIN	RM-0.5+	Jones	Personal observation: both juvenile COH and SH were found.	1986
	SH	RM-0.5+	Brown and Moyle Mitchell & Coyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Harris	Electrofished: 2 sites, juvenile COH and SH were found 6,523 feet above the mouth. Carcass survey: found CHIN, COH and SH present.	1993 1996/97
North Fork Redwood Creek T21N, R16W, Sec. 17	SH	RM-?		A stream survey is needed.	
Muddy Gulch Creek T21N, R16W, Sec. 16	?	RM-?		A stream survey is needed.	
Kenny Creek T21N, R16W, Sec. 22	SH	RM-0.5	Jones	Personal observation: juvenile SH were found at the crossing, 0.5 miles upstream above the mouth. CHIN have been seen in this stream by others.	1986
	COH	RM-?			
	CHIN	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
Mud Creek T21N, R16W, Sec. 26	SH	RM-?	Fender, Litwin and Swirhun	Survey: none observed because of the muddy water. A local landowner reported that "salmon" used to spawn there but he hasn't seen any for a few years.	1980
			Jones	Personal observation: juvenile SH were found near the mud springs. The extent of their stream use was unknown.	1986

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Bear Creek T21N, R16W, Sec. 25	CHIN	RM?	Sprehn & Clark	Survey: 1 CHIN were found above the mouth during this February survey.	1979
	COH	RM-?	Jones	Personal observation: juvenile SH were present in the stream.	1991
	SH	RM-1.0	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1986
Taylor Creek T21N, R16W, Sec. 36	SH	RM-0.25	Michaels and Anderson	Survey: juvenile SH in densities of 75/ 100 feet of stream were present. RT (resident fish) inhabited the headwater (0.7 mi.).	1969
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
	RT	RM-0.7			
Little Rock Creek T21N, R15W, Sec 32	SH	RM- ?	Jones	Personal observation: SH were see above the fishway under the Branscomb Road.	1986, 1997
Section Four Creek T21N, R15W, Sec. 33	SH	RM-1.4	Jones	Personal observation: juvenile SH were seen in this stream.	1979
Hulls Creek (tributary to main stem Eel River. This portion in Mendocino Co. Only T25N, R13W, Sec. 33-1/2 at Co. line)	SH	RM- 13.8 (Mendocino Co. Only)	Barney	Personal observation: caught SH from the Pepperwood Creek tributary down to the mouth.	1953/54
	CHIN	RM-?	Cherr & Griffin Jones	Fisheries inventory: SH and CHIN were present. Personal observations: a major stream system with very little fishery information. Spot checks from time to time revealed the presence of SH and CHIN. This stream needs a comprehensive survey of the main stem and tributaries.	1979 1965-1997
Horse Canyon Creek T25N, R13W, Sec. 33-1/2	?	RM-?		Appears to be a significant stream with no information on the fishery. A stream survey is needed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Lost Cabin Creek T24N, R13W, Sec. 27-1/2	?	RM-?		Appears to be a significant stream with no information on the fishery. A stream survey is needed.	
Who Who Creek T24N, R13W, Sec. 13-1/2	?	RM-0.25+	Barney	Personal observation: fished the lower 1/4 mile of stream catching juvenile SH. Appears to be a significant stream with no fishery information. A stream survey is needed.	1968-72
Hulls Valley Creek T24N, R13W, Sec 20-1/2	SH	RM-6.5 +	Mullins, Humphrey, Goodfield & Downey	Electrofished: 3 sites, found juvenile SH 34,471 feet upstream. This appears to be a significant stream and needs further surveys.	1995
Pepper Wood Creek T24N, R 12W, Sec. 21	?	RM-?		Appears to be a significant stream with no fishery information. A stream survey is needed.	
Cardenas Creek T24N, R12W, Sec. 11	?	RM-?	Barney Jones	Personal observation: fished juvenile SH in this stream in the early 1960's. SH have access upstream to the falls located 3/4 mile above the mouth. This stream appears to be significant but very little fishery information is available. It needs to be surveyed.	Early 1960's 1997
Chamise Creek (in Mendocino Co.) T4S, R6E, Sec. 33	SH	RM-?	Yoshioka	Personal observation: juvenile SH were present in 1999 where the Alder Point road crosses the stream. Appears to be a significant stream and needs to be surveyed.	1999
Charlton Creek T5S, R5E, Sec. 26	?	RM-?		Appears to be a significant stream with no information on the fishery. It needs to be surveyed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
North Fork Eel River (This portion in Mendocino County) T24N, R14W, Sec. 4	SH	RM- 10 (in Mendo. Co.)	Carfel & LaFaunce	Survey: from Hulls Creek To Salt Creek, juvenile SH were most abundant in pools, (Jim Gilman of CDFG) reported SSH present in the headwater in his youth.	1973
	SSH	RM-?			
	CHIN	RM-10	Holman & Baker	Survey: many juvenile SH were seen in July.	1973
				Finlayson & Alberson	Survey: No salmonids were seen in September, high temperatures was suspected - a barrier at split rock stopped anadromous migrations at the time. This barrier was removed in 1986 under a CDFG grant.
			Walton and Gilroy	Carcass survey: found no fish.	1988
			Jones	Personal observation: Native Americans trapped CHIN and SH in a traditional location near Split Rock on the North Fork Eel River. Chicken wire fences, spears and hook and line were the primary methods of take. The native Americans in the area also report SSH present, however, none were confirmed during a survey by E. Gurstung and Jones in 1985,a year before the barrier was removed. What was found was late run winter SH in the process of spawning.	1997
Horse Creek T24N, R14W, Sec. 13	SH	RM-0.5+	Jones	Personal observations: juvenile SH were seen up to RM-0.5 miles. The full extent of use was unknown. A full stream survey is recommended.	1997
Asbill Creek T24N, R13W, Sec. 60	SH	RM-0.5	Jones	Personal observations: juvenile SH and CHIN carcasses were seen. This stream has a good supply of cool summer flow. It is often fished by Native Americans. Habitat restoration work has been done under contract between the Tribe and CDF&G.	1995, 1996, 1997
	CHIN	RM-?	Smith & Bevitori	Survey: juvenile SH were seen 1,584 feet upstream.	1996
West Fork Asbill Creek	SH	RM- 0.5	Jones	Personal observation: Juvenile SH were present. This stream has a good source of cool flow.	1995
Bear Canyon Creek T24N, R13W, Sec. 53	SH?	RM-?		This stream needs to be surveyed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Bell Springs Creek, (tributary to main stem Eel River) T24N, R14W, Sec. 18	SH	RM0.4	Jones Goodrich & Melton	Personal observation: juvenile SH were present, but the extent of their range was unknown. Electrofished: 1 site, juvenile SH were found 2,037 feet upstream.	1995 1996
Blue Rock Creek T24N, R14W, Sec. 30	SH	RM-2.0	Jones	Personal observation: Juvenile SH were present.	1989
Shell Rock Creek T23N, R14W, Sec. 17	SH	RM-2.5	Cherr & Griffin Jones	Fisheries inventory: SH use this stream. Personal observation: Juvenile SH were present during the summer.	1979 1981
Corbet Creek T23N, R14W, Sec. 33	?	RM-?		This stream needs to be surveyed.	
Black Oak Creek T23N, R14W, Sec. 33	?	RM-?		This stream needs to be surveyed.	
Gill Creek T23N, R14W, Sec. 2	SH	RM-3.0	Moore	Field note: juvenile SH were present above the section line between Section 6 and 7.	1981
Woodman Creek T22N, R14W, Sec. 11	SH SSH CHIN	RM-5.9 RM-2.4 RM-2.4+	Jones & Swirhun Jones Brown & Bevitori	Survey: the mouth of the stream was once a barrier, but CDF&G constructed a fishway over it. Summer steelhead once occupied the stream, (Merv Pinches) but no longer are present (Jones). CHIN also used the stream, but their current range is unknown. Personal observations: juvenile SH were present, the fishway at the mouth needs frequent maintenance. Survey: juvenile SH were found 15,399 feet upstream.	1981 1988 1995
Iron Creek T22N, R14W, Sec. 16	SH	RM-3.3	Jones	Personal observations: Juvenile steelhead were found 3.3 miles upstream. A survey of the entire stream is needed.	1981

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
White Rock Creek T22N, R14W, Sec. 9	SH	RM-2.6	Jones	Personal observations: juvenile steelhead were present during spot checks. The stream needs to be surveyed in its entirety.	1981
Board Tree Canyon T22n, R14W, Sec. 18	SH	RM-1.0	Jones	Personal observations: juvenile SH were seen. A more comprehensive survey is needed.	1981
Stoney Creek T22N, R14W, Sec. 24	?	RM-?		A stream survey is needed.	
Burger Creek T22N, R14W, Sec. 36	SH CHIN	RM-6.6 RM-0.3	Cherr & Griffin Jones Jones Brown & Bevitori	Fisheries inventory: SH use this stream. Personal observations: found only juvenile SH present. Personal observation: the rock barrier once present 0.3 miles up has been removed and SH have been permitted to migrate upstream. CHIN have been seen below the old barrier. Juvenile SH were present 22,636 feet upstream, (4.3 miles) and, at times seen 6.6 miles up stream.	1979 1984 1991 1995
One Eyed Creek T22N, R14W, Sec. 35	SH	RM-1.0	Jones	Personal observation: spot check revealed the presence of juvenile SH and/or RT. A survey is needed.	1991
Bear Pen Canyon T22N, R14W, Sec. 10	SH	RM-3.5	Shapovalov & Vestal Jones Brown & Bevitori	Survey: This stream in Bear Pen Canyon was known as Cold Creek in 1938. Juvenile SH were present. Personal observations: spot checks revealed the presence of juvenile SH and/or RT. Survey: juvenile SH were seen 18,441 feet upstream.	1938 1991 1995

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Middle Fork Eel River T21N, R13W, Sec. 6	SH	RM-47.3 (in Mendo. Co.)	Jones	<p>Personal observation: Including annual surveys to determine the population of SSH (varied from an adult population of 198 in 1966 to 1,601 in 1981). The 2000 count was 279 adult fish, well below the 500 recommended by the Threatened Trout Committee as the minimum necessary to maintain the resource and the point where the fish should be listed as threatened or endangered.</p> <p>A management plan has been prepared by the US Forest Service and CDFG. Winter run SH are believed to occupy the same range as the SSH, but evidence is lacking. CHIN cannot migrate above the roughs at Osborne. Spring run CHIN were observed during two survey years, but now appear depleted. The SSH and possibly SH runs in the stream extend into the North Fork of the Middle Fork and Balm of Gilead Creek tributaries in Trinity County. In the main stem they were once found upstream to Uhl Creek where a barrier falls exists but a barrier at the lower end of Wrights Valley prevents current upstream access beyond this point.</p> <p>Above the barrier at Wrights Valley a RT population was present up to Willow Basin Creek in the headwater.</p>	1966-2000
	SSH	RM-47.3 (in Mendo. Co.)			
	CHIN	RM-38.0			
	COH	RM-?			
	RT	RM-10.5+			
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. (COH are not present in the area above the U.S. Forest Station on Hwy 162. They may not be present in this drainage at all. W. Jones).	1991
Goforth Creek T21N, R13W, Sec. 6	SH	RM-2.0	Ashuckian & Pinkham	Survey: found low flow, poor spawning and nursery area. Juvenile SH in low numbers occupy the stream up to the roughs 200 to 300 yards above mouth. A falls exists 2 miles upstream.	1976
Eastman Creek T21N, R13W, Sec. 5	?	RM-?		A stream survey is needed.	
Salt Creek T21N, R13W, Sec. 11	SH	RM-4.5	Jones & Pinkham	Survey: spot checked, Juvenile SH were present in the lower area in numbers of 300/100 feet of stream. There may be resident trout in the headwater. A more extensive survey is recommended.	1976

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Elk Creek T21N, R11W, Sec. 18	SH CHIN	RM-14.8 RM-?	Leos Walton and Griffin Jones and Flosi Goodfield, MacDonnell & Mesman Jones Daugherty	Field note: CHIN carcasses were found. Carcass survey: the lower two miles was surveyed, no fish were found. Carcass survey: lower mile, found one CHIN carcass. .Survey: found juvenile SH present 12,822 feet above the mouth. Personal observations: observation were limited to 2 miles above the mouth with juvenile SH present. This is a significant stream and a more extensive survey is needed. Observation: Found juvenile SH present at 14.8 miles, (T20N, R11W, Sec 36).	1983 1987/88 1987/88 1994 1995 1995
Deep Hole Creek T20N, R11W, Sec. 5	?	RM-?		A stream survey is needed.	
Shake Creek T20N, R11W, Sec. 5	?	RM?		A stream survey is needed.	
Sanhedron Creek T20N, R11W, Sec. 5	SH/RT	RM-?	Daugherty	Observation: Found juvenile salmonids (SH/RT) in June.	1995
Bennett Creek T20N, R11W, Sec. 9	?	RM-?		A stream survey is needed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Little Thatcher Creek T20N, R11W, Sec. 4	?	RM-?		A stream survey is needed.	
Bear Creek T20N, R11W, Sec. 23	?	RM-?		A stream survey is needed.	
Crocker Creek T20, R11W, Sec. 26	SH/RT	RM4.3	Daugherty	Snorkeled: found juvenile salmonids in July, 4.3 miles upstream (T19N, R11W, Sec. 4).	1995
Sulphur Springs Creek T20N, R11W, Sec. 25	SH? RT Brook trout	RM-3 RM-1.0+ RM-1.0+	CDFG Daugherty Jones & Daugherty	The CDF&G resource map indicates a possible falls near the mouth. Snorkel: June 29, Temperature 11 ⁰ C, found juvenile salmonids (SH/RT) and brook trout present (T19N, R11W, Sec 10). However, only (SH/RT) found at the lower site (T19N, R11W, Sec. 35.). Personal observations: Brook trout were found in “Cow Glade” in the headwaters. Electrofishing revealed 27 brooktrout and one RT. A stream survey of the entire drainage is needed.	1979 1995 1996
Mendenhall Creek T20N, R11W, Sec. 36	?	RM?		A stream survey is needed.	
Sportsman Creek T20N, R11W, Sec. 36	SH	RM-0.09	Daugherty	Observation: Found juvenile SH present in September about 500 feet above the mouth.	1995
Thatcher Creek T21N, R11W, Sec. 7	SH CHIN	RM-6.8 RM-4.5	Kershner Jones	Survey: stream survey by BLM, SH were present, primarily in the upper areas. Erosion and sedimentation limit fishery use. Personal observation: SH and CHIN use this stream. Further surveys are needed.	1974 1995

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Haynes Creek T21N, R11W, Sec. 14	?	RM-?		This is a significant stream and an extensive survey is needed.	
Skunk Lake Creek T21N, R11W, Sec. 11	?	RM-?		A stream survey is needed.	
Red Rock Creek T21N, R11W, Sec. 11	?	RM-?		A stream survey is needed.	
Hanson Creek T21N, R11W, Sec. 11	?	RM-?		A stream survey is needed.	
Jack Hollow Creek T21N, R11W, Sec. 14	?	RM-?		A stream survey is needed.	
Hayshed Creek T21N, R11W, Sec. 8	?	RM-?		A stream survey is needed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Mill Creek T22N, R12W, Sec. 25	SH CHIN COH	RM-15 RM-11.7+ RM-?	Jones Leos Brown and Moyle Goodfield, Melton and Ouradnik CDF and USFS	Personal observations: including review of CDFG grant proposals and carcass surveys. Both SH and CHIN have been observed. Carcass survey: 2 CHIN carcasses were found downstream of the Adobe lane bridge. Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. Electrofished: found juvenile SH present 35,376 feet upstream. Carcass survey: no fish were found.	1958-97 1983 1991 1996 1998/99 and 1999/00
Turner Creek T22N, R12W, Sec. 22	SH CHIN	RM-4.0 RM-?	Jones	Personal observation: including review of CDFG grant proposals and spot checks at road crossings. Both SH and CHIN have been seen. A restoration project including the construction of a fishway has been completed on this stream.	1985-1997
Unnamed stream (South Fork Turner Creek) T22N, R12, Sec. 20	SH	RM-?	Jones	Personal observation: includes review of CDFG grant proposals, juvenile SH were seen.	1995
Grist Creek T22N, R12W, Sec. 9	SH COH	RM-? RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. A stream survey is needed.	1991
Town Creek T22N, R12W, Sec. 7	SH	RM-3.0+	Barney Jones	Personal observation: juvenile SH were seen above the old defunct reservoir, near the confluence of Doan Creek. Personal observation: and spot checks, most of the Round Valley section of the stream was either of intermittent flow or was dry during the late spring and summer. Juvenile SH have been see in the upper portion of the stream on the west side of the valley. However, the extent of their use was not determined. A stream survey is needed.	Early 1950's 1966-00

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Doan Creek T22N, R13W, Sec. 3	?	RM-?		A stream survey is needed.	
Little Valley Creek T22N,R13W, Sec. 2	?	RM-?		A stream survey is needed.	
Doan Creek T22N, R13W, Sec. 3	?	RM-?		A stream survey is needed.	
Short Creek T22N, R12W, Sec. 4	SH CHIN	RM-8.25 RM-?	Jones Jones Goodfield, Milton & Ouradnik	Personal observation: carcass surveys, spot checks and review of CDFG grant projects: both CHIN and SH were seen in this stream. Carcass surveys: found CHIN present. Electrofished: 1 site, found juvenile SH 16,836 feet upstream.	1966-1997 1987/88 and 1988/89 1996
Tank Creek T23N, R16, Sec. 16	SH	RM-?	Jones	Personal observation: including spot checks, juvenile SH were seen.	1995
Alder Creek T23N, R13W, Sec. 15	SH	RM-0.8	Jones	Survey: juvenile SH were seen up to the 20 foot barrier falls.	1995
Mud Creek T23N, R13W, Sec. 15	SH	RM-0.5	Jones	Personal observation: juvenile SH were seen up to the 15 foot falls.	1995
Cold Creek T23N, R13W, Sec. 15	SH RT	RM-2.2 RM-2.7+	Jones	Personal observation: juvenile SH were seen up to the 30 foot falls. A resident RT population was present above.	1995

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Silver Creek T23N, R12W, Sec. 36	?	RM-?		A stream survey is needed.	
Williams Creek T23N, R11W, Sec. 31	SH	RM-3.9	Baracco & Johnson	Survey: SH were present from the barrier at Blueschist Narrows down.	1976
	CHIN	RM-?	Reynolds	Carcass survey: saw 7 live chinook and 2 carcasses in Section 13 and 24.	1981
			Leos	Carcass survey: found 3 CHIN carcasses.	1983
			Jones	Personal observation: this stream has been fished by native Americans using chicken wire netting placed in stream, bank to bank. CHIN and SH populations are currently low. A resident trout population was believed present above Blueschist Narrows.	1966-1997
			Goodrich & Melton	Survey: found juvenile SH present 9,801 feet upstream.	1996
			Daugherty	Carcass survey: Daugherty of NMFS observed a CHIN speared on a redd in this stream.	1999/00
Poor Mans Creek T23N, R12W, Sec. 25	CHIN	RM-1.0	Jones	Personal observation: when rebuilding Hwy 162 in the Williams Creek Canyon a problem developed with CHIN passage at the Poor Mans Creek culvert. Numerous CHIN were unable to pass the culvert. The Federal HWY project team corrected the problem to permit passage.	1985
Murphy Creek T23N, R12W, Sec. 13	SH	RM-3.0	Jones	Personal observation: spot checks and CDFG grant proposal application reviews were conducted. Juvenile SH were seen in this stream.	1995
	CHIN	RM-1.3	Reynolds	Carcass survey: 10 live CHIN and 2 carcasses were found.	1981

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Black Butte River T23N, R11W, Sec. 28	SH CHIN	RM-9.5 in Mendo. Co., 5.6+ in Glenn Co. RM-?	Thompson & Jones Jones Smith	Survey: Both SH and CHIN use this stream. In the 1930's this stream had SSH. Juvenile SH were seen during the 1970 survey. The SH fishery extends 5.6+ miles into Glenn County. Electrofishing: found juvenile SH present below the road crossing in the headwater (Basin area). Carcass survey: found CHIN present.	1970 1986 1999-00
Van Horn Creek T23N, R11W, Sec. 27	?	RM-?	Jones	Personal observation: the mouth of the stream dries up during the summer. A stream survey is needed.	1970
Shields Creek T23N, R11W, Sec. 34	?	RM-?	Jones	Personal observation: the mouth of the stream becomes dry during the summer. A stream survey is needed.	1970
Nebo Creek T23N, R11W, Sec. 36	?	RM-?	Jones	Personal observation: The mouth of the stream becomes dry during the summer. A stream survey is needed.	1997
Jump off Creek T23N, R11W, Sec. 36	SH	RM-3.9	Coleman & Van Zandt	Survey: juvenile SH were considered abundant, 25/100 feet of stream.	1975
Billy Williams Creek T22N, R11W, Sec. 1	?	RM-?	Jones	Personal observation: The mouth of the stream becomes dry during the summer. A stream survey is needed.	1970
Sallady Creek T22N, R10W, Sec. 7	?	RM-?	Jones	Personal observation: The mouth of the stream becomes dry during the summer. A stream survey is needed.	1970
Mississippi Creek T22N, R10W, Sec. 16	?	RM-?	Jones	Personal observation: The mouth of the stream becomes dry during the summer. A stream survey is needed.	1970

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
White Hawk Creek T22N, R10W, Sec. 16	?	RM-?	Jones	Personal observation: The mouth of the stream becomes dry during the summer. A stream survey is needed.	1970
Blue Slides Creek T22N, R10W, Sec. 15	?	RM-?	Jones	Personal observation: The mouth of the stream becomes dry during the summer. A stream survey is needed.	1970
Atchison Creek T22N, R10W, Sec. 26	?	RM-?		Heads in Glenn County. A stream survey is needed.	
Pinto Creek T22N, R10W, Sec. 26	?	RM-?		This stream heads in Glenn County. A stream survey is needed.	
Billy Pike Creek T22N, R10W, Sec. 26	?	RM-?	Jones	Personal observation: the mouth of this stream becomes dry during the summer. A stream survey is needed.	1970
Butte Creek T22N, R10W, Sec. 35	?	RM-?	Jones	Personal observation: the mouth of this stream becomes dry during the summer. A stream survey is needed.	1970
Dingman Creek T22N, R10W, Sec. 35	?	RM-?	Jones	Personal observation: The mouth of this stream becomes dry during the summer. A stream survey is needed.	1970
Salmon Creek T21N, R10W, Sec. 1	?	RM-?	Jones	Personal observation: The mouth of this stream becomes dry during the summer. A stream survey is needed.	1970

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Cold Creek T21N, R10W, Sec. 12	SH	RM-0.75	Burger	Survey: a barrier, 10 feet high was found 3/4 mile above the mouth. Juvenile SH number 250-300/100 lineal feet in the lower stream. Above the barrier was a resident RT population The stream had a good summer flow and cool water. During the survey the flow was 6 CFS at the mouth and water temperature was 48-52 ^o F in October.	1973
	RT	RM-8.5	Jones	Electrofished: the first pass in a 100 foot section of stream resulted in the capture of 32 juvenile SH/RT. The second pass resulted in the capture of 26 juvenile SH/RT. The stream heads in Glenn County to the east of Mendocino County.	1981
			Daugherty	Electrofishing: found RT present 8.5 miles upstream of the falls (within Glenn Co.)	1995
Cottonwood Creek T21N, R8W, Sec. 7	RT	RM-0.5	Daugherty	Personal observation: found RT present 0.5 miles above the mouth, (Glenn Co.).	1995
Baldy Creek T21N, R10W, Sec. 24	SH	RM-0.25	Burger	Survey: juvenile SH were present in the lower 1/4 mile of stream, a grade change of 100 feet/ 100 yards acts as a barrier. No resident trout were seen above.	1973
Buckhorn Creek T21N, R10W, Sec. 24	SH	RM-3/4	Burger	Survey: juvenile SH were present in numbers of 35-60/100 linear feet of stream. Above the falls was a resident population, range unknown.	1973
	RT	RM-?			
Rocky Basin Creek T21N, R9, Sec. 19	?	RM-?		The stream heads in Glenn County. A stream survey is needed.	
Hunter Creek T21N, R 9W, Sec. 29	?	RM-?		Needs to be surveyed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Skidmore Creek T21N, R9W, Sec. 32	SH	RM-0.25	Burger	Survey: juvenile SH were seen below the barrier at 1/4 mile. Up stream, juvenile SH were found with a population numbering 15 fish /100 linear feet of stream. The confluence of this stream with the Black Butte River is in Glenn County, with the headwater in Mendocino County.	1973
French Creek T21N, R9W, Sec. 32	?	RM-?		Needs to be surveyed.	
Umbrella Creek T21N, R9W, Sec. 32	?	RM-?	Jones	Personal observation: the confluence of this stream with the Black Butte River is in Glenn County. However, it's headwater is in Mendocino County. A stream survey is needed.	1970
White Horse Creek T20N, R9W, Sec 4	?	RM-?		Needs to be surveyed	
Spanish Creek T20N, R9W, Sec. 10	RT	RM-3.7	Daugherty	Electrofished: found RT present 3.7 miles upstream in Glenn Co.	1995
Basin Creek T20N, R9W, Sec. 15	?			In Glenn Co. Needs to be surveyed.	
Estell Creek T20N, R9W, Sec. 15	?			In Glenn Co. Needs to be surveyed.	
Middle Creek T20N, R9W, Sec. 15	?			In Glenn Co. Needs to be surveyed.	
Bar Creek T23N, R11W, Sec. 14	SH	RM-0.3	Jones O'Leary	Personal observation: spot checks revealed a small juvenile SH/ RT population. Survey: found resident RT present from above Road M1 to the river, a distance of 1,800 feet.	1991 1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Buck Creek T23N, R11W, Sec. 11	?	RM-?	Jones Shea and Richie	Personal observations: the mouth of this stream is dry during the summer, (as seen during Middle Fork summer steelhead surveys conducted annually 1966-1999). Survey: no fish were observed during the survey from the M1 road to the mouth.	1966-1999 1997
Travelers Home Creek T23N, R11W, Sec. 2	SH	RM-0.2	Jones Shea and Richards	Personal observation: a boulder roughs at high grade restricts SH from moving out of the Middle Fork inner canyon. However, the lower 0.2 milrd of this stream within the Middle Fork canyon supports juvenile SH in numbers of 50/ 100 linear feet of stream. Survey: no fish were observed in the headwaters. Juvenile SH were found in the lower 800 feet of stream.	1997 1997
Fly Creek T24N, R10W, Sec. 31	RT	RM-Barrier at mouth	Cassidy, Campman and Mulchaey	Survey: resident trout were present throughout the drainage, however a steep bouldered section at the mouth prevents anadromous entry.	1995
Devils Den Creek T24N, R10W, Sec. 31	SH?	RM-?	Jones	Personal observation: the lower end of the stream near the Middle Fork Eel dries up each summer. A survey is needed.	1966-1999
Howard Creek T24N, R10W, Sec. 30	RT	RM-?	Priest, Piovarcsik	Survey: an 8 foot falls and a steep gradient prevent anadromous access. However, rainbow trout are present throughout the drainage above.	1995
Bear Creek T24N, R10W, Sec. 30	none	RM-?	Jones Sullivan and Crand	Survey: the stream gradient was steep and the mouth of this stream has been dry every summer. No fish were observed.	1966-1999 1995
Fish Town Creek T24N, R10, W, Sec. 19	SH	RM- 50 yards	Jones	Personal observation: yearly- 1966-1999, upstream access out of the Middle Fork Eel River inner canyon has been blocked by an extensive steep grade.	1966-1999

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Beaver Creek T24N, R10W, Sec. 17	SH SSH RT	RM-0.3 RM-0.3 RM-5.5	Harris	Survey: observed juvenile SH and an adult SSH in the lower 0.3 miles of the stream. Further migration has been blocked by a barrier falls and steep grade. A resident population of coastal rainbow resides above.	1995
Hammerhorn Creek T24N, R10W, Sec. 18	RT	RM- 3.4	Jones Cassidy and Priest	Personal observation: a barrier falls was present at the mouth. A resident population of coastal rainbow use the stream above the falls. Survey: the resident RT population uses 3.4 miles of stream.	1966-1999 1995
Smokehouse Creek T24N, R10W, Sec. 15	RT	RM-2.0	Harris and Ritchie	Survey: found RT present in the stream up to a barrier falls of 5 feet in height about ½ mile above Forest road M1.	1995
Pothole Creek T24N, R10W, Sec. 12	SH	RM-1.5	Richards and Shea	Survey: juvenile SH were present 1.5 miles upstream.	1997
Fossil Creek T24N, R11W, Sec. 12	SH	RM-0.3	Jones O'Leary	Personal observation: annual observation during the Middle Fork Eel summer steelhead surveys revealed juvenile SH in the lower reach. The population extends to a falls, a series of boulder and bedrock at a grade change of (30-60 %). Survey: no fish were seen above the falls.	1966-1999 1997
Alder Creek T24N, R11W, Sec. 1	none	RM-0	Jones O'Leary and Richards	Personal observation: the mouth of this stream has been seen yearly between 1966-1997. A grade change barrier was present at the mouth. Above the mouth was a series of steep roughs as the stream courses through the Middle Fork Eel River (MFE) inner canyon. Survey: no fish were seen.	1966-1999 1997
Maple Creek T24N, R11W, Sec. 1	none	RM-0	Jones Shea	Personal observation: Within the Middle Fork Eel River Canyon was a series of continuous steep roughs considered to be barriers. Survey: no fish were seen.	1966-1999 1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Deer Lick Creek T24N, R11W, Sec. 2	none	RM-0	Jones Richards and Shea	Personal observation: Within the Middle Fork Eel River Canyon was a series of barrier roughs. This stream heads in Trinity County. Survey: no fish were seen.	1966-1999 1997
Rattlesnake Creek T25N, R11W, Sec. 35	SH RT	RM-50 yards RM-3.0	Hicks , McCurdy Brown and Moyle Ritchie and Shea	Survey: juvenile SH were present in the lower 50 yards but were blocked from further upstream access by high gradient roughs. Resident rainbow trout were present above the roughs, up to 3 miles above the mouth. This stream heads in Trinity County. Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. (Highly doubted, saw none during Middle Fork Eel River surveys 1966-1997 . W Jones). Survey: RT were found upstream 4,840 feet.	1966 1991 1997
Stick Lake Canyon Creek T25N, R11W, Sec. 34	none	RM-0	Jones O'Leary and Nishijuma	Personal observation: Within the Middle Fork Eel River Canyon was a series of continuous steep gradient barriers. Survey: no fish were found.	1966-1999 1997
Line Gulch T25N, R11W, Sec. 34	none	RM-0	Jones	Personal observation: the mouth of this stream is in Trinity County, yet the major portion is in Mendocino County. The mouth of this stream was seen yearly between 1966 and 1997 No fish were present. A grade change forms a barrier at the mouth.	1966-1999
See Trinity County for additional, headwater Middle Fork Eel River Tributaries)					
Mainstem Eel River tributaries (upstream of the Middle Fork Eel River)					

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Dean Creek T21N, R13W, Sec. 7	SH	RM-0.7	Coyle & Crittenden	Electrofished: 3 sites, found juvenile SH present 3,585 feet upstream.	1992
Brad Turner Creek T21N, R13W, Sec. 18	?	RM?		This stream needs to be surveyed.	
Division Creek T21N, R13W, Sec. 19	none	RM-0	Jones	Personal observation: a barrier falls occurs at the mouth.	1995
Rock Creek T21N, R13W, Sec. 19	SH	RM-0.1	Jones Mullins & Humphrey	Personal observation: there was a barrier falls near the mouth. Survey: juvenile SH were present up to the falls 532 feet above the mouth.	1995 1996
Tatu Creek T21N, R13W, Sec. 19	?	RM-?		This stream needs to be surveyed.	
Rodeo Creek T21N, R13W, Sec. 30	?	RM-?		A stream survey is needed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Outlet Creek T21N, R13W, Sec. 31	SH	RM-18.0	Durkee & Klamt	Survey: juvenile SH were present. Spawning and rearing areas were mainly in the tributaries and mid and lower main stem.	1972
	COH	RM-16.0	Leos	Carcass survey: 11 CHIN carcasses and 9 skeletons were seen.	1983
	CHIN	RM-18.0	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones	Personal observation: CDFG grant projects and an adult trapping project located near the mouth of Long Valley Creek has taken place on this stream. The trapping project was for the purpose of obtaining CHIN eggs. During the trapping project many adult CHIN were caught for egg taking to rear and release young salmon in an effort to evaluate ocean catch. One chum salmon was caught and released during the trapping operations. COH use the mid drainage and a few tributaries.	1985, 1990-1997
			Jones	Tissue samples were taken: tissues were taken from 12 male CHIN and 4 female CHIN and given to Dr. Jennifer Nielsen for genetic work..	1995
			Goodfield & John	Electrofished: 2 sites, found juvenile SH present 60 feet downstream of the Willits sewer treatment plant.	1995
			Harris	Carcass survey: found CHIN present.	1998/99
			Harris	Snorkel survey in July. Found juvenile CHIN and SH present.	1999
			Harris	Carcass survey: CHIN were present.	1999/00
Bloody Run Creek T20N, R14W, Sec. 1	SH	RM-2.6	Jones	Personal observation: juvenile SH were present.	1975
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Darr & Terwilliger	Personal observation: juvenile SH were found upstream 12,429 feet.	1995
			CDF and USFS	Carcass survey: CHIN were present.	1998/99 and 1999/00

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Cherry Creek T20N, R15W, Sec. 2	SH	RM-1.5	Grass	Field Note: a barrier rock falls blocks anadromous upstream migration 1.5 miles above the mouth. Junenile SH were present downstream.	1986
			Terwilliger, John & Darr	Survey: juvenile SH were found up to a 13 foot high dam, 7,847 feet above the mouth of the stream.	1995
Long Valley Creek T20N, R14W, Sec. 28	SH	RM-5.1	Durkee & Klamt	Survey: juvenile SH were present.	1972
	COH	RM-?	Reynolds	Carcass survey: 230 live CHIN were counted. A total of 35 carcasses and 2 skeletons were seen.	1981
	CHIN	RM-5.0	Leos	Carcass survey: 84 CHIN carcasses and 9 skeletons were found.	1983
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Goodrich, John & Short	Electrofished: found juvenile SH present at 26,849 feet upstream.	1995
			Harris	Carcass survey: found CHIN present, but no COH.	1998/99 and 1999/00
Sam Watt Creek T20N, R14W, Sec. 29	?	RM-?	Harris	Carcass survey: no fish were present in the lower 1/4 mile. The stream needs to be surveyed.	1999/00
Dutch Henry Creek T20N, R14W, Sec. 29	SH	RM-1.5	Durkee & Klamt	Survey: Juvenile SH were present in the lower stream. A 50 foot barrier falls was found 1.5 miles above the mouth.	1972
	COH	RM-?	Long & Griffin	Carcass survey: 7 CHIN carcasses were found.	1987
	CHIN	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Harris	Carcass survey: no fish were seen.	1999/00

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Reeves Canyon Creek T19N, R14W, Sec. 11	COH	RM-1.5	Jones	Carcass survey: Found COH and CHIN present in the stream.	1987
	SH	RM-1.5	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
	CHIN	RM-1.5			
Rowes Creek T19N, R14W, Sec. 23	RT	RM-?	Jones	Personal observation: a barrier roughs was found near the mouth. Resident trout were present above the barrier in low numbers. The stream needs to be surveyed.	1990
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
Ryan Creek T19N, R14W, Sec. 26	SH	RM-1.0	Durkee & Klamt	Survey: found juvenile SH present.	1972
	CHIN	RM-1.0	Jones	Carcass survey: 126 adult CHIN and 17 redds were found. Three barrier culverts block further upstream migrations of anadromous fish.	1985
	RT	RM-0.5+	Jones	Carcass survey: in 1987 a carcass survey revealed 7 CHIN and 20 redds.	1987
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones	Electrofishing: started in 1986 and has been conducted annually since. COH were present through 1993. RT were present above the three barrier culverts but the extent of their population remains unknown.	1986-1999
			Harris	Electrofished: no COH were present, but juvenile SH were present in 1997, 1998, and 1999.	1997-98-99
Harris	Carcass surveys: found CHIN present both seasons.	1998/99 and 1999/00			
Bull Creek T19N, R14W, Sec. 26	?	RM-?	Harris	Personal observation: the HWY 101 culvert was a problem. The stream needs to be surveyed.	2000

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Berry Creek T19N, R13W, Sec. 31	SH	RM-5.0	Jones	Personal observation: juvenile SH were present in the canyon. The Little Lake Valley portion of the stream becomes dry each summer.	1997
			Harris	Personal observations: no fish have been seen above the valley floor.	2000
Davis Creek T19N, R13W, Sec. 31	SH	RM-6.0	Lincoln & Wong	Electrofished: below Morris Dam, a barrier, the surveyors found juvenile SH in a density of 54 fish/ 100 feet of stream.	1974
	CHIN	RM-6.0	Coleman & Van Zandt	Stream survey: estimated 60 juvenile SH were present/ 100 feet of stream. The stream throughout the upper end of Little Lake Valley becomes dry during the summer.	1975
			Harris	Personal observation: a spot check off the bridge at the south east end of the valley revealed one adult SH.	1999/00
Moore Creek T18N, R13W, Sec. 28	?	RM-?		This stream needs to be surveyed.	
Conklin Creek T18N, R13W, Sec. 33	SH	RM-0.2	Jones	Survey: juvenile SH present were present below the barrier - boulder roughs located 0.2 miles above the mouth. And, resident trout were present in 2.0 miles of stream above.	1997
	RT	RM-2.0			
Mill Creek T19N, R13W, Sec. 31	CHIN	RM-4.0	Harris	Carcass survey: found CHIN and SH present.	1998/99 and 1999/00 1991
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	
	SH	RM-4.0	Harris	Electrofishing: two stations, found juvenile SH present in both.	1998 and 1999
Upp Creek T18N, R13W, Sec. 7	CHIN	RM-0.5	Jones	Personal observation: one CHIN carcass was observed below the HWY 101 crossing. However, the range of this species remains unknown. A stream survey is needed.	1985

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³	
Willits Creek T18N, R14W, Sec. 13	SH	RM-3.0	Klamt & Edwards	Survey: SH were present.	1971	
	CHIN	RM-3.0	Reynolds	Carcass survey: 97 adult CHIN were observed.	1981	
	COH	RM-?	Leos	Carcass survey: six surveys found a total of 99 CHIN carcasses, and 32 skeletons.	1983	
	RT	RM-0.5+		Reynolds	Carcass survey: 97 carcasses and 8 skeletons were found.	1985
				Jones & Reynolds	Carcass survey: 308 CHIN and 58 skeletons were found.	1986
				Walton, Gilroy, Griffin and Ferrara	Carcass survey: found 363 CHIN and one COH. Further fish movement upstream was blocked by a reservoir.	1987
				Jones	Carcass survey: found 144 CHIN and one COH present in Willits Creek.	1988
				Jones	Carcass survey: found 106 CHIN and no COH	1989
				Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
				Goodrich & John	Electrofishing: 1 site, juvenile SH were found 5,564 feet upstream of the mouth. Carcass surveys: found CHIN present.	1995
				Jones and Harris		1996/97
				Harris	Carcass surveys: CHIN and SH were present in 1997/98. CHIN were also seen during the carcass surveys of 1998/99 and 1999/00.	1997/98, 1998/99 and 1999/00
				Harris	Downstream migrant trapping: was conducted and found both juvenile CHIN and SH present.	1999 and 2000
NMFS				Observation: found juvenile COH present during the summer of 2001	2001	
Dutch Henry Creek T18N, R14W, Sec. 2	RT	RM-1.0	CDFG	Resource map: this stream is above a dam and reservoir operated by the Brooktrails Resort Improvement District. The dam was constructed in the 1960's. What was once SH habitat is now RT habitat.	1979	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³	
Baechtel Creek T18N, R13W, Sec. 6	SH	RM-9.0	Jones	Survey: juvenile SH were present, CHIN were also known to use the stream.	1980	
	CHIN	RM-8.0	Jones	Personal observation: CDFG grant program to remove debris barriers to downstream migrant CHIN and SH in the floor of the valley was successful. Debris jams and increases in quantities of in-stream gravels and debris forced the stream to crest banks and flood the valley floor to a depth of from 1-3 inches to upwards of 8 feet. This forced the salmonids on to the valley floor where water temperatures were in the mid 80's. Many of the downstream migrants attempting to migrate downstream through the flat, shallow, high temperature valley floor perished and were lost. At the time of CDFG intervention CHIN and COH had expired from all but Mill Creek. Trapping was required to move the juvenile CHIN around the blocked areas to maintain the fishery resource during this critical period. The grant provided funds to remove the debris barriers and excess gravel, reducing the stream bed to grade and allowing the anadromous salmonids to again re-use most of the valley streams. One portion of the debris extraction included the removal of 1,000 automobile and truck tires.	1986	
	COH	COH-8.0				
				Walton	Carcass survey: found 4 SH, 19 CHIN and 1 COH. Local residents indicated many more were present prior to the timing of the CDFG survey.	1987
				Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
				Terwilliger & Darr	Survey: found juvenile SH present 36,925 feet, (7.0 miles) upstream.	1995
			Jones	Personal observations: COH have not been found in Bechtel Creek in recent years.	1997	
			Harris	Carcass surveys: CHIN and SH were found during the surveys.	1998/99 and 1999/00	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³	
Haehl Creek T18N, R13W, Sec. 13	SH	RM-4.0	Durkee & Klamt	Survey: reported the stream to be of limited value to anadromous fish.	1972	
	CHIN	RM-4.0	Berg	Electrofishing: fish rescue activities found juvenile steelhead present.	1988	
	COH	RM-4.0	Jones	Brown and Moyle	Carcass survey: CHIN were seen. In the Jan 26, survey 75 CHIN and 1 COH were found, on the Feb 1, survey 20 CHIN and 5 COH were found.	1987
					Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
					Electrofishing: 2 sites, juvenile SH were found 7,922 feet upstream.	1995
					Carcass surveys: revealed no fish either season.	1998/99 and 1999/00
Broaddus Creek T18N, R13W, Sec. 13	COH	RM-4.5	Walton	Carcass survey: found 11 COH present.	1987	
	SH	RM-4.5+	Jones	Personal observation: have seen CHIN, COH and SH present in this stream.	1985-1999	
	CHIN	RM-2.0+	Jones	Brown and Moyle	Electrofished: at the HWY 20 Cal/trans bridge west of Willits and found juvenile COH and SH present. Doubt that any COH currently (2000) use this stream.	1990
					Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
					Electrofished: 1 site, juvenile SH were found 720 feet upstream.	1995
					Personal observations: spot checks revealed redds. Fish were not seen.	1998/99
Harris		Harris	Carcass surveys: found CHIN present.	1999/00		
Alder Creek T20N, R13W, Sec. 5	?	RM-?		A stream survey is needed.		
Brushy Creek T20N, R13W, Sec.4	?	RM-?		A stream survey is needed.		

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Indian Creek T20N, R13W, Sec. 9	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. A stream survey is needed.	1991
Twin Bridges Creek T19N, R12W, Sec. 6	?	RM-?		Needs to be surveyed.	
Salt Creek T19N, R12W, Sec. 20	?	RM-?		A stream survey is needed.	
Garcia Creek T19N, R12W, Sec. 22	SH	RM-2.0	Burger	Survey: Juvenile SH were present in the upper area in a density of 40 fish/100 feet of stream.	1973
Thomas Creek T19N, R12W, Sec. 26	SH	RM-1.4	Burger	Survey: juvenile SH were present in densities of 15-50/100 feet of stream.	1973
Tomki Creek T18N, R12W, Sec. 12	SH CHIN	RM-18.0 RM-18.0	Durkee & Klamt Jones Goodfield, Miller & Schaibler Harris	Survey: Found SH present. Carcass surveys: SH and CHIN were present. The Soil Conservation Service CHIN population estimates for the years 64/65 through 87/88 were: 64/65 1,747, 65-75 no survey, 75/76 367, 76/79 no survey, 79/80 2,362, 80/81 317, 81/82 565, 82/83 1,741, 83/84 no survey, 84/85 1,292, 85/86 3,558, 86/87 2,173, 87/88 3,611. Electrofished: juvenile SH were present near the mouth of Cave Creek.	1972 1964-1988 1997
Salmon Creek T18N, R12W, Sec. 15	SH CHIN	RM-? RM-?	Jones	Personal observation: carcass surveys revealed CHIN and juvenile SH present. A stream survey is needed.	1985

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Long Branch Creek T18N, R12W, Sec. 9	SH CHIN	RM-2.5 RM-2.5	Durkee & Klamt Jones Harris	Survey: juvenile SH were present. Carcass survey: found 79 CHIN in the stream, 28% of the total for Tomki Creek this year. Carcass surveys: conducted in 1997/98, 1998/99 and 1999/00 found CHIN present each year.	1972 1984 1997/98, 1998/99 and 1999/00
Scott Creek T18N, R12W, Sec. 8	?	RM-?		A stream survey is needed.	
Cave Creek T18N, R12W, Sec. 18	SH CHIN	RM-2.7 RM-?	Jones Harris	Carcass survey: juvenile SH and CHIN were seen, alive and as carcasses. The county road, a secondary road to State HWY 101, fords this stream 6 times and produces an appreciable amount of silt and sediment. These stream "ford" crossings should be bridged. Carcass surveys: conducted in 1997/98, 1998/99 and 1999/00, found CHIN and a few SH present each year.	1984-1997 1997/98, 1998/99 and 1999/00
Unnamed Tributary to Cave Creek T18N, R112W, Sec. 18	?	RM-?		A stream survey is needed.	
Little Cave Creek T18N, R12W, Sec. 19	?	RM-?		A stream survey is needed.	
Unnamed Tributary to Tomki Creek T18N, R13W, Sec. 12	?	RM-?		A stream survey is needed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
String Creek T19N, R13W, Sec. 36	SH	RM-5.5	Durkee & Klamt	Survey: juvenile SH were present.	1972
	CHIN	RM-4.1	Jones	Carcass survey: numerous surveys found CHIN present annually.	1984-1997
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			CCC Harris	Survey: juvenile SH were seen up to 21,612 feet upstream. Carcass surveys: conducted in 1997/98, 1998/99 and 1999/00, found CHIN present each year.	1995 1997/98, 1998/99 and 1999/00
Digger Creek T19N, R13W, Sec. 35	?	RM-?		A stream survey is needed.	
Rocktree Creek T19N, R13W, Sec. 35	SH	RM-1.8	Patton	Personal observation: juvenile SH and CHIN were present.	1982
	CHIN	RM-1.8	Jones	Personal observation: SH and CHIN use this stream. Summer flow was low.	1984-1997
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Harris	Carcass survey: CHIN were found.	1998/99
Tartar Creek T19N, R13W, Sec. 26	SH	RM-1.2	Durkee & Klamt	Survey: juvenile SH were present.	1972
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
Wheelbarrow Creek T19N, R13W, Sec. 9	SH	RM-1.7	Jones	Personal observation: and carcass surveys, SH and CHIN were present.	1984-1997
	CHIN	RM-1.7	Harris	Carcass surveys: conducted in 1997/98, 1998/99 and 1999/00, found CHIN present each year.	1997/98, 1998/99 and 1999/00
Whitney Creek T18N, R11W, Sec. 18	SH	RM-2.0	Burger	Survey: juvenile SH were present, in densities of 15 fish /100 feet in the lower reach and 50/100 feet of stream in the headwater.	1973

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Hale Creek T18N, R11W, Sec. 19	?	RM-?		A stream survey is needed.	
Salmon Creek T18N, R11W, Sec. 30	?	RM-?		A stream survey is needed.	
Mill Creek T18N, R11W, Sec. 30	?	RM-?		A stream survey is needed.	
Trout Creek T18N, 11W, Sec. 34	SH RT	RM-0 RM-?	Grass	Personal observation: a boulder barrier at the mouth prevents access. However a resident trout population extends upstream an unknown distance.	2000
Bucknell Creek T18N, 11W, Sec. 35	SH	RM-0.2 in Mendocino Co. Additional in Lake Co.	Hann Grass	Survey: juvenile SH were seen. Survey: juvenile SH seen, abundant, 20-40 fish /100 feet of stream.	1959 1990
See Lake County for mainstem Eel River and tributaries for additional data					
Headwater of the Eel River above Lake County					
Hull Creek T20N, R10W, Sec. 25	RT	RM-2.1	Reese & Abel	Survey: found RT present in numbers of 15/ 100 feet of stream. A barrier, located about ½ mile below Forest road 20N16, prevented further upstream access. This main stem Eel River tributary is entirely within Mendocino County.	1993
Wesley Ives Creek T20N, R10W, Sec. 25	RT	RM-0.9	Reese & Abel	Survey: RT were present in numbers of about 3 fish/100 feet of stream. This stream is entirely within Mendocino County.	1993

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³	
Russian River (Mendocino County portion) (see Sonoma County for the lower portion of the drainage)	COH	RM-25.8	Jones	Personal observation: There is a vast amount of information on the fishery resources of the Russian River, but will not dwell upon this here other than to refer readers to two documents: (1) "The Russian River Drainage--A Summary Report on the Fish and Wildlife Resources and their Problems, By the Inland Fisheries and Wildlife Staffs of Region 3, Calif. Department of Fish and Game, 1969", and (2) "A History of the Salmonid Decline in the Russian River, Prepared by: Steiner Environmental Consulting, 1996".	1958-1999	
	SH	RM-25.8				
	CHIN	RM-25.8				
				Jones	COH in recent years have been infrequently present in the upper drainage (Mendocino County). In the late 1980's a substantial number migrated into the upper river. Again in the 1999-00 season Bill Townsend caught one in the Ukiah area, (verified by a CDFG Warden). However these fish only reach the upper river during certain years and may be limited to periods, when flows are sufficient for fish passage at the Healdsburg Recreational Dam.	1985-1999
				Jones	SH are the most numerous of the three salmonid species using the upper river. Egg taking operations occur at the mitigation hatchery located at the base of the Coyote Valley Dam. Records are available from CDFG.	1958-1999
			Jones	Electrofishing: 6 juvenile CHIN, and numerous juvenile SH were caught while looking for tule perch in the river upstream of Hopland in May.	1987	
			Jones	CHIN numbers in this section of river have been low and have varied from year to year. During the fall 1999 season anglers reported many in the upper river. Their occurrence in the upper river may also be limited by low flow years.	1999	
			Jones	Memo: report to Steiner Environmental Consulting, 3 CHIN were seen on a one day (a 7 mile float) on the upper main stem.	1999	
Ash Creek T12N, R11W, Sec. 36	SH RT	RM-1.4 RM-0.8	Jones	Survey: RT were found in the headwater above the SH habitat. A number of substantial barrier falls, some as great as 25 feet were present in the mid area. The trout population is limited by low stream flow.	1997	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Edwards Creek T12N, R11W, Sec. 25	SH	RM-2.3	Reed & Gillespie Jones & Mizuno	Survey: juvenile SH were found to use the stream up 2.3 miles. Survey: found juvenile SH in the lower section of stream, 3 redds were also found, they were presumed to be from SH that spawned the previous season.	1976 1990
Cummiskey Creek T12N, R11W, Sec. 14	SH	RM-7.0	Vestal	CDF&G report: Russian River Drainage - A Summary Report. Indicates 7.0 mile of fishery habitat.	1969
McDonald Creek T12N, R11W, Sec. 9	?	RM-?		A stream survey is needed.	
Pieta Creek T12N, R11W, Sec. 2	SH	RM-10.0	Bruns & Jackson P.G.E. Sommarstrom	Survey: juvenile SH were present in substantial numbers, 35/100 feet in the lower area and in numbers of 150-200 fish/ 100 feet of stream in the headwater near the confluence of Tyler Creek. Electrofished: numerous stations, the juvenile SH population was estimated at 22,700 fish. Electrofished: 5 sites, the juvenile SH population was estimated at 46,700 fish.	1974 1976 1985
Coleman Creek T12N, R11W, Sec. 2	SH	RM-6.2	P.G.E. Sommarstrom	Electrofished: near the Younce Road crossing, found juvenile SH. The projected population of juvenile SH for Coleman Creek was 17,450 fish. Electrofished: 5 sites, the juvenile SH population was estimated at 31,500 fish.	1976 1985
Salt Canyon Creek T12N, R11W, Sec. 1	?	RM-?		A stream survey is needed.	
Vasser Creek T13N, R11W, Sec. 35	SH	RM-?	CDFG	Resource map: SH present. A stream survey is needed.	1979

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Jakes Creek T13N, R10W, Sec. 32	?	RM-?		A stream survey is needed.	
Salt Springs Creek T12N, R10W, Sec. 7	?	RM-?		A stream survey is needed.	
Tyler Creek T12N, R10W, Sec. 15	SH	RM-6.0	Johnson	Survey: juvenile SH were present up to the instream reservoir found upstream of the Tyler Valley Road crossing. The densities of juvenile SH found in Tyler Creek below the reservoir was about 225 fish/ 100 feet of stream.	1974
			P.G.E.	Electrofished: found juvenile SH present up to the reservoir. The estimated population of RT in this stream was 24,430 fish.	1976
			Sommarstrom	Electrofished: 4 sites, found juvenile SH present. Their population estimate was 34,300 fish.	1985
Hoil Creek T12N, R9W, Sec. 18	SH	RM-1.6	P.G.E.	Electrofishing: found juvenile SH present in Hoil Creek up to 1.6 miles. The juvenile SH population for the stream was estimated to be 2,300 fish.	1976
			Sommarstrom	Electrofishing: 5 sites, the estimated total population of juvenile SH was 6,040 fish.	1985
Feliz Creek T13N, R11W, Sec. 19	SH CHIN	RM-13.5 RM-?	Meda	Watershed restoration report: a review of the application included a reference to the catching of a CHIN in 1974.	1988
			Jones	Personal observation: walked the stream observing juvenile SH and non game fish.	1985-1997
Duncan Creek T13N, R12W, Sec. 23	?	RM-?		A stream survey is needed.	
Johnson Creek T13N, R12W, Sec. 26	?	RM-?		A stream survey is needed.	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Young Creek T13N, R12W, Sec. 7	?	RM-?		A stream survey is needed.	
Middle Fork Feliz Creek T13N, R12W, Sec. 18	?	RM-?		A stream survey is needed.	
North Fork Feliz Creek T14N, R13W, Sec. 36	?	RM-?		A stream survey is needed.	
Dooley Creek T13N, R11W, Sec. 19	SH	RM-2.8	Jones Lester, Jenkins, Canata, and Coey	Personal observation: worked on water applications and road projects in this drainage. SH use this system for spawning and juveniles are present during the summer months. Electrofished: found juvenile SH present at 14,679 feet above the mouth of the stream. A culvert 8 feet in diameter, 80 feet long (possibly a barrier) was present at 15,335 feet upstream of the mouth.	1964-1997 1998
North Fork Dooley Creek T13N, R11W, Sec. 15	SH	RM-0.3	Lester, Jenkins, Canata, and Coey	Electrofished: found juvenile SH at two locations, 747 feet upstream and 1,752 feet upstream.	1998
Middle Fork Dooley Creek T13N, R11W, Sec. 22	SH	RM-0.5	Lester, Jenkins, Canata and Coey	Electrofished: found juvenile SH present at 3 locations. The uppermost station was at 2,619 feet above the mouth.	1998
McDowell Creek T13N, R11W, Sec. 22	SH	RM-3.2	CDFG CDFG Lester, Jenkins, Canata and Coey	Observation: “ several 0+ age sized RT/SH were present in the ravine at the foot of the Hopland Grade”, July 8, 1954. CDF&G report: 50 juvenile SH were observed over 75 yards of stream on June 5th. Electrofished: found 70 juvenile at 17,118 feet above the Russian River. A 15 foot falls was present at 20,071 feet above the mouth of the stream.	1954 1990 1998

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
McNab Creek T14N, R12W, Sec. 36	SH	RM-1.5	CDFG Parloto, T. and M. Miller	CDFG report: fish rescue records on McNab Creek in 1955, 56, 58, 59, 60, 61, 62, 63, 64, and 65 include catching juvenile SH. Electrofished: found juvenile SH up to a dam, not thought to be a barrier, at 7,953 feet above the mouth. Discussions with the land owner, "Fuzzy" indicate that SH may migrate up to and 1/4 mile beyond the reservoir at 23,053 feet. Summer flows were very low.	1955-65 1997
Parsons Creek T14N, R12W, Sec. 25	SH	RM-3.6	CDFG Miller, M., B. Kieffer, and C. Adams Coey Opperman, J.	CDFG report: fish rescue records for the years 1949, 50, 55, the 1960's, 70, 71, 72, and 1973 include catching juvenile SH. Electrofishing: found juvenile SH present, "the end of anadromy was the cascading waterfall at 19,177 feet above the mouth". Carcass survey: observed one adult SH in a pool above the ford at the UC. Davis, Hopland Field Station. Juvenile SH were rescued during a period of decreasing flows from a location below the ford in July. Summer stream flows were low. Personal observation: juvenile SH were present in intermittent pools below a point 19,177 feet above the mouth of the stream.	1949-73 1998 1999 1999
Morrison Creek T14N, R12W, Sec. 15	?	RM-?		A stream survey is needed.	
Howell Creek T14N, R12W, Sec. 3	?	RM-?		A stream survey is needed.	
Robinson Creek T14N, R12W, Sec. 4	SH	RM-5.4	CDFG Huber, A., J. Fort and K. Morgon	CDFG report: in ". . . 1954, abundant RT & SH were observed". "In August of 1956, report a 100 foot section of the stream yielded 223 young SH." Juvenile SH were found in the upper middle section in August of 1962. In July of 1973, 25 juvenile SH were present in an 100 foot section of stream above the gorge. In the gorge the population was 100 juvenile SH per 100 feet of stream. Electrofished: found juvenile SH at 28,299 feet above the mouth, (access above this point was not granted by the landowner).	1954, 56, 62, 73 1995

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Unnamed tributary to Robinson Creek T14N, R12W, Sec. 7	?	RM-?		A stream survey is needed.	
South Branch Robinson Creek T14N, R13W, Sec. 11	SH	RM-2.9	Jones	Personal information: hiked the stream looking at possible restoration projects. Found juvenile SH present 2.9 miles.	2000
Unnamed Tributary to Robinson Creek T14N, R13W, Sec. 11	SH	RM-?	CDFG Morgon, K. and K. Gregory	CDFG report: records show that Robinson Creek was planted with steelhead from Prairie Creek in 1938, from Crawford Creek, Dooley Creek, Feliz Creek, Johnson Creek, McNab Creek, and other areas of Robinson Creek in 1958. Juvenile SH were also planted in Robinson Creek from McNab Creek in 1960, 61, 62, 63, 66, 72, and in 1975 from Talmage ponds. Electrofished: found juvenile SH present 3,202 feet above the mouth.	1938-75 1995
Skunk Creek T15N, R13W, Sec. 34	?	RM-?		A stream survey is needed.	
Mill Creek T15N, R12W, Sec. 28	SH RT	RM-3.0 RM-1.0	Elwell Jones	Survey: the stream above the upper Talmage pond supports a resident population of RT. SH utilize the stream below. Personal information: anadromous SH utilize the stream below the lower Talmage pond and tributaries for spawning and rearing purposes. CDFG plants catchable trout in the two impoundments on this stream annually.	1958 1958-1997
McClure Creek T15N, R12W, Sec. 27	SH	RM-2.7	Baracco	Survey: while the lower end of the stream becomes dry during the summer, the upper portion of the stream provides habitat for juvenile SH up to the 20 foot falls.	1976
North Fork Mill Creek T15N, R12 W, Sec. 26	SH RT	RM-1.5 RM-1.0	Finlayson and Nelson	Survey: SH utilize the lower 1.5 miles of stream for spawning and rearing. A 10 foot falls blocks passage and was considered a barrier. A resident trout population was present between the lower falls and a 50 foot high falls located one mile further up the stream.	1973

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Doolin Creek T15N, R12W, Sec. 28	SH	RM-2.3+	Jones	Personal observation: juvenile SH have been seen in this stream.	1995
Gibson Creek T15N, R12W, Sec. 28	SH RT	RM-1.5 RM-2.5+	Jones	Personal observation: SH were able to migrate upstream into the City of Ukiah, however a culvert barrier under the Leslie Street crossing stops further access. The fish were blocked from spawning and rearing above. Upstream of this barrier was a resident trout population. At one time in the past the CDFG operated a small hatchery on this stream, located west of the town at the foot of the mountain. CDFG also permitted the planting of adult SH surplus to the Coyote Valley facility, in this stream for spawning purposes. The extent of the headwater population of RT is unknown.	1985-1997
Orrs Creek T15N, R12W, Sec. 16	SH RT	RM-6.0 RM-4.0	Jones Jones	Survey: Orrs Creek offers roughly six miles of steelhead spawning and nursery area. About 2.5 miles upstream is a bedrock boulder area that currently is the subject of habitat restoration work. Personal observation: a CDFG grant program is expected to work on this stream in 2000. When anadromous fish cannot migrate above the barrier site resident trout utilize the area above. CDFG has permitted the planting of adult SH surplus to the needs of the Coyote Valley Hatchery, in the headwater of this stream enabling them to benefit from the better spawning and rearing habitat.	1990-1997 1999
Sulphur Creek T15N, R12W, Sec. 16	SH RT	RM-0.8 RM-2.5+	Jones	Personal observation: SH ascend Sulphur Creek to a falls of 8 feet in height. Above this falls is a small resident trout population in the mountainous terrain heading south east of the Ukiah valley. Within the RT area are three additional barriers: one a 10 foot high on-stream dam located 1.0 miles upstream of the first barrier falls, two, a bedrock falls of 15-20 feet high located one mile further upstream and three, a 20 foot on-stream dam is located an additional ½ mile upstream.	1985-97

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Ackerman Creek T15N, R12W, Sec. 8	SH	RM-10	CDFG	CDFG report: the lower 3.5 miles of stream was chemically treated in 1954 and partially again in 1965.	1974
			Jackson and Bruns	Survey: juvenile SH were present for 10 miles of stream.	1974
			Swirhun and Litwin	Survey: the streambed stabilization structure at North State Street was considered a barrier as was the culvert crossing at 6.0 miles out the Masonite road. The streambed stabilization structure has since been repaired.	1980
			Swirhun and Ambrosius	Field note: juvenile SH were seen all the way up the stream indicating the streambed stabilization structure and culvert were not stopping fish migrations at this time.	1981
			Jones	Personal observation: the fish ladder (streambed stabilization structure) under North State Street was rebuilt in 1983 and has functioned since.	1983-1997
			Koehler, C., S. Shapleigh and L. Gray	Electrofished: found juvenile SH present at 18,524 feet above the mouth.	1994
Hensley Creek T15N, R12W, Sec. 5	SH	RM-3.75	Holman	Survey: the HWY 101 crossing was assumed to be a barrier. Below this were juvenile SH.	1957
			Bruns	Survey: no barriers were observed, juvenile SH were present throughout 3.75 miles of stream. The juvenile SH abundance was 25-30 per 100 feet of stream.	1974
			Jones, Sanders and Swirhun	Survey: found juvenile SH present in numbers of 200/100 feet of stream in the mid section and 50 /100 feet in the lower area. A log barrier was present 1 mile above the narrow canyon west of the valley floor.	1980
			Jones	Personal observation: CDFG memo, an inspection was conducted of a newly completed fishway under HWY 101. The fishway was found to be functioning properly.	1987
Howard Creek T15N, R12W, Sec. 8	?	RM-?	Jones	Personal observation: this stream was intermittent at best and serves as the spillway for the Coyote Valley Dam.	1997

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
East Fork Russian River T15N, R12W, Sec. 5	SH RT COH	RM-1.0 RM-10.0 RM-?	Jones Brown and Moyle	Personal observation: Coyote Valley Dam (Lake Mendocino) was constructed on this stream in 1957. The fisheries management program in the reservoir above is based on a warm water fishery. CDFG currently operates a mitigation type salmonid egg taking facility at the foot of this dam. The program is funded by the Corps of Engineers. Records of fish captured, spawned and released are available from the CDFG. On occasion, CHIN have been captured at the facility. Resident rainbow trout are present in the drainage above the reservoir. The CDFG plants catchable trout in the canyon area above the reservoir. The CDFG has also planted brown trout in the canyon above the reservoir. Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1958-2000 1991
Cold Creek T16N, R11W, Sec. 18	RT	RM-2.0+	Jones	Personal observation: much of this stream parallels HWY 20 and is visible from the road. Resident rainbow trout use this stream although a major portion dries up during the summer months. At one time CDFG maintained a small hatchery on this stream.	1964-1997
Mewhinney Creek T16N, R11W, Sec. 5	RT	RM-?		A stream survey is needed.	
White Creek T16N, R11W, Sec. 5	RT	RM-?	Denasion	Personal observation: resident RT were present on a seasonal basis. This stream needs to be surveyed.	2000
Burright Creek T17N, R11W, Sec. 32	RT	RM-3.0+	Lee and Namba Jones	Survey: found RT upstream to 3.0 miles. Personal information: RT have been seen 3.0 miles upstream. Additional surveys are needed.	1978 1985
Bevans Creek T17N, R11W, Sec. 20	?	RM-?	Jones	Personal observation: an instream dam and reservoir were located 1.8 miles above the mouth. The fishery resources in the stream and reservoir are unknown. A stream survey is needed.	1997
Williams Creek T17N, R11W, Sec. 17	?	RM-?		A stream survey is needed.	

Stream ¹/Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Busch Creek T17N, R11W, Sec. 8	RT	RM-5.0+	Jones	Personal observations: RT have been seen 5.0 miles upstream. The upper limit of habitat for the RT is unknown. The stream needs to be surveyed.	1984-1986
Boyes Creek T17N, R11W, Sec. 12	?	RM-?		A stream survey is needed.	
Adobe Creek T17N, R11W, Sec. 7	?	RM-?		A stream survey is needed.	
Power House Canal T17N, R11W, Sec. 7	RT	RM-1.3	Jones	Personal observation: this section of stream contains RT (some of the salmonids enter the stream through the trans basin diversion from the Eel River). This diversion is now screened from October until June.	1997-2000
York Creek T16N, R12W, Sec. 32	SH COH	RM-3.0 RM-?	Netherby and Rockwood Jones Brown and Moyle	Survey: low summer flows, juvenile SH were seen. The survey team did not continue on the main stem but turned up the unnamed stream (South Fork). Fishery resources above the South Fork were not surveyed. Personal observation: juvenile SH were present in pools in the main stem above the (South Fork for another 1.5 miles). Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1966 1966 1991
Unnamed tributary to York Creek T16N, R12W, Sec. 29	SH	RM-0.5	Netherby and Rockwood	Survey: juvenile SH were found up to the falls 0.5 miles above the mouth.	1966

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Forsythe Creek T16N, R12W, Sec. 16.	SH	RM-7.5	Brackett	Survey: juvenile SH in numbers of 150-200 fish/100 feet of stream, up to the barrier at RM-7.5. Above this barrier RT were present for an additional 5.0 miles.	1963
	CHIN	RM-4.0+	Durkee and Klamt	Survey: the survey noted the bedrock/boulder barrier restricting anadromous access and also a second barrier 5.0 miles upstream in Section 8, limiting further RT access. Additional limitations include low flow and high summer stream temperatures.	1972
	COH	RM-?	Jones	Personal observation: some of the earlier Fishery Biologists that preceded me in the central coastal area have indicated that COH once used some of the tributaries to this drainage. A CHIN carcass was found on a stream bar above the HWY 101 crossing.	1958-1965
	RT	RM-5.0	Brown and Moyle Freele, B. and S. Carey	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. Electrofished: found juvenile SH present at 15,509 feet above the mouth.	1991 1999
Seward Creek T16N, R12W, Sec. 7	SH	RM-2.0	Brackett	Survey: in his survey of Forsythe Creek Brackett indicated that local people spoke about "silver salmon" having been present in the drainage at one time.	1963
	COH	RM-2.0	Durkee and Klamt	Survey: juvenile SH were present.	1972
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones	Personal observation: Fishery Biologists present in the central coastal during the late 1950's and 1960's indicated that COH once used this drainage. However, I have never seen any in this stream.	1997
		Freele	Electrofished: Juvenile SH were present 3,097 feet above the mouth.	1999	

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Eldridge Creek T16N, R13W, Sec. 11	SH COH	RM-5.1 RM-?	Brackett Brown and Moyle Jones Higgins, and Freele	Survey: found juvenile SH. In his survey of Forsythe Creek Brackett indicates that local people spoke of "silver salmon" having been present in the drainage at one time. Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. Personal observation: conducted CDFG grant application reviews in the drainage and have also heard from Fishery Biologists in the 1950's and 1960's indicated that COH were once present in this stream. The barriers that were present in the 1970's were removed in the 1980's . However, COH are no longer present. Electrofished: found juvenile SH present 26,820 feet upstream.	1963 1991 1997 1999
Jack Smith Creek T16N, R13W, Sec. 11	SH COH	RM--5.5 RM-?	Brackett Durkee and Klamt Brown and Moyle Jones	Survey: found juvenile SH. In Brackett's survey of Forsythe Creek he indicates that local people spoke of "silver salmon" having been present in the drainage at one time. Survey: full of barriers to anadromous fish, only resident trout are present. Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. Personal observation: conducted CDFG grant application reviews in the drainage and have also heard from Fishery Biologists that were present in the 1950's and 1960's have indicated that COH were once present in this stream. The barriers that were present in the 1970's were removed in the 1980's . However, COH are no longer present.	1963 1972 1991 1997
Baker Creek T16N, R12W, Sec. 6	SH	RM-1.4	Freele, Higgins, and Jankowski	Electrofished: 5 sites, found juvenile SH present in the stream as high as 7,353 feet.	1999

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Mill Creek T17N, R13W, Sec. 36	SH	RM-4.2	Brackett	Survey: In Brackett's survey of the Forsythe Creek Drainage he states, "It appears that silver salmon utilize this drainage to a limited extent." He further said, "It appears that Mill Creek has great importance to the Forsythe Creek drainage as a spawning and nursery area".	1963
	RT	RM-3.5			
	COH	RM-?	Durkee and Klamt	Survey: the lower 4.5 miles of drainage was open to SH and juveniles were present. A boulder barrier blocks further access. "An additional 3.5 miles of drainage is available with removal of the barrier."	1972
			Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
			Jones	Personal observation: Fishery Biologists present in the 1950's and 1960's have indicated that COH were once present in this stream. However, I have never seen any in this stream. In the last few years the Ukiah Rod and Gun have with CDFG approval, planted surplus adult SH from the Coyote egg taking station into this stream at a location near RM 31,080.	1997
		Freele, B. and S. Higgins	Electrofished: juvenile SH were found at 21,392 feet above the mouth.	1999	
		Freele, B. and A. Fairbrook	Electrofished: "SH" were present at 41,340 feet (7.8 miles) above the mouth. The survey indicated SH however they were probably RT as the barrier remains in place. (W. Jones)	1999	
Walker Creek T17N, R13W, Sec. 21	RT	RM-1.0	Jones and Irwin	Survey: found RT below and above the reservoir. The trout above the reservoir were limited from further access by a bedrock falls in Section 12. The RT population above the reservoir was about 50 fish/100 feet of stream. Below the reservoir RT have access to about 1.0 miles of stream and population levels were low.	1978
			Freele, B. and H. Sean	Electrofished: 4 sites, juvenile SH were found as high as 3,731 feet above the mouth of the stream.	1999
Salt Hollow Creek T16N, R12W, Sec. 9	SH	RM-0.4	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
	COH	RM-?	Mullen and Harris	Stream survey: found juvenile SH present in low numbers in Salt Hollow Creek 0.4 miles above the mouth on July 14th. Suspect they migrated further upstream before a Dam was constructed.	2000

Stream ¹ /Tributary (RM=river mile)	Species / Run	Upper Limit of Run ² (RM=river mile)	Sources References / Pers. comm.	Comments	Survey Dates ³
Rocky Creek T16N, R12W, Sec. 20	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991. A stream survey is needed.	1991
Mariposa Creek T17N, R12W, Sec. 17	SH	RM-0.5	Rockwood	Survey: an instream dam and reservoir was present and limits SH to 0.5 miles of access. A resident trout population extends 1.2 miles above the reservoir.	1966
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
	RT	RM-1.2	Jones	Personal observation: the RT population continues to exist. The current landowner has expressed interest in restoring the SH use to the upper drainage.	2000
Fisher Creek T17N, R12W, Sec. 17	SH	RM-1.8	Rockwood	Survey: juvenile SH were present. "It was an excellent spawning and nursery stream".	1966
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991
Corral Creek T17N, R12W, Sec.	SH	RM-1.6	Jones	Personal observation: juvenile SH were present in the stream up 1.6 miles. A stream survey is needed.	1966
	COH	RM-?	Brown and Moyle	Report: COH were listed in this stream in the report; "Status of Coho Salmon in California", 1991.	1991